

Lie n -groupoids and stacky Lie groupoids

Chenchang Zhu *
 Departement Mathematik
 ETH Zentrum, Rämistrasse 101
 8092 Zürich, Switzerland
 (zhu@math.ethz.ch)

February 2, 2008

Abstract

We discuss two generalizations of Lie groupoids. One consists of Lie n -groupoids defined as simplicial manifolds with trivial $\pi_{k \geq n+1}$. The other consists of stacky Lie groupoids $\mathcal{G} \Rightarrow M$ with $\mathcal{G}$ a differentiable stack. We build a 1-1 correspondence between Lie 2-groupoids and stacky Lie groupoids up to a certain Morita equivalence. Equivalences of these higher groupoids are also described.

Contents

1	Introduction	2
2	Lie n-groupoids	6
2.1	The description of Lie 2-group(oid)s with finite data	7
3	SLie groupoids	11
3.1	Good charts	13
4	From SLie-groupoids to Lie 2-groupoids	17
4.1	The construction of $X_2 \rightrightarrows X_1 \rightrightarrows X_0$	17
4.2	Proof that what we construct is a Lie 2-groupoid	19
5	From Lie 2-groupoids to stacky Lie groupoids	21
5.1	The stack $\mathcal{G}$	21
5.2	Proof that $\mathcal{G} \Rightarrow M$ is an SLie groupoid	21
6	Equivalences of Lie 2-groupoids	25
6.1	Strict maps of Lie homotopy types	25
6.2	Morita Equivalence of Lie 2-groupoids	28
6.3	1-Morita Equivalence of Lie 2-groupoids	32

*Research partially supported by the Liftoff fellowship 2004 of the Clay Institute

7 One-to-one correspondence between Lie 2-groupoids and SLie groupoids 32

8 Comments on the inverse map 36

1 Introduction

Recently there has been much interest in higher group(oid)s, which generalize the notion of group(oid)s in various ways. Some of them turn out to be unavoidable to study problems in differential geometry. An example comes from the string group, which is a 3-connected cover of $Spin(n)$. More generally, to any compact simply connected group G one can associate its string group $String_G$. It has various models, given by Solz and Teichner [Sto96] [ST04] using an infinite dimensional extension of G , by Brylinski [BM94] using a $U(1)$ -gerbe with the connection over G , and recently by Baez et al. [BCSS] using Lie 2-groups and Lie 2-algebras. Henriques [Hen] constructs the string group as a higher group in the setting of this paper and as an integration object of a certain Lie 2-algebra using an integration procedure similar to that of [Š, Zhu].

Other examples come from a kind of étale stacky groupoids (Weinstein groupoids) [TZ06a]. They are the 1-1 global objects integrating Lie algebroids, where a Lie algebroid could be roughly understood as a bundle of Lie algebras. Notice that unlike (finite dimensional) Lie algebras which always have associated Lie groups, Lie algebroids do not always have associated Lie groupoids [AM84, AM85]. One needs to enter the world of stacky groupoids to obtain the desired 1-1 correspondence. Since Lie algebroids are closely related to Poisson geometry, this result applies to complete the first step of Weinstein's program of quantization of Poisson manifolds: to associate to Poisson manifolds their symplectic groupoids [Wei83, WX91]. It turns out that some "non-integrable" Poisson manifolds cannot have symplectic (Lie) groupoids. This problem is solved in [TZ06b] with the above result so that every Poisson manifold has a corresponding étale stacky symplectic groupoid.

Higher group(oid)s were already studied in the early twentieth century by Whitehead and his followers under various terms, such as cross-modules. However in this paper we use a uniform method to describe higher (respectively Lie) groupoids using simplicial sets [May92] (respectively manifolds) since it is believed that there should be an equivalence between n -groupoids and spaces whose homotopy groups are trivial above π_n (also called n -coskeleta or n -truncated homotopy types)¹. The 0-simplices of the simplicial set correspond to the objects, the 1-simplices to the arrows (or 1-morphisms), and the higher dimensional simplices the higher morphisms. This method becomes more suitable especially when we are dealing with the differentiable category.

Recall that a simplicial set (respectively manifold) X is made up by sets (respectively manifolds) X_n and structure maps

$$d_i^n : X_n \rightarrow X_{n-1} \text{ (face maps)} \quad s_i^n : X_n \rightarrow X_{n+1} \text{ (degeneracy maps)}, \text{ for } i \in \{0, 1, 2, \dots, n\}$$

¹This was suggested (indirectly) by Jacob Lurie to the author, however it was known much earlier, for example, by Duskin and Glenn [Dus79, Gle82].


that satisfy the coherence conditions

$$\begin{aligned} d_i^{n-1} d_j^n &= d_{j-1}^{n-1} d_i^n \text{ if } i < j, & s_i^n s_j^{n-1} &= s_{j+1}^n s_i^{n-1} \text{ if } i \leq j, \\ d_i^n s_j^{n-1} &= s_{j-1}^{n-2} d_i^{n-1} \text{ if } i < j, & d_j^n s_j^{n-1} &= \text{id} = d_{j+1}^n s_j^{n-1}, & d_i^n s_j^{n-1} &= s_j^{n-2} d_{i-1}^{n-1} \text{ if } i > j+1. \end{aligned} \quad (1)$$


The first two examples are the simplicial m -simplex $\Delta[m]$ and the horn $\Lambda[m, j]$ with

$$\begin{aligned} (\Delta[m])_n &= \{f : (0, 1, \dots, n) \rightarrow (0, 1, \dots, m) | f(i) \leq f(j), \forall i \leq j\}, \\ (\Lambda[m, j])_n &= \{f \in (\Delta[m])_n | \{0, \dots, j-1, j+1, \dots, m\} \not\subseteq \{f(0), \dots, f(n)\}\}. \end{aligned} \quad (2)$$


In fact the horn $\Lambda[m, j]$ is a simplicial set obtained from the simplicial m -simplex $\Delta[m]$ by taking away its unique non-degenerate m -simplex as well as the j -th of its $m+1$ non-degenerate $(m-1)$ -simplices, as in the following picture (in this paper all the arrows are oriented from bigger numbers to smaller numbers):



A simplicial set X is *Kan* if any map from the horn $\Lambda[m, j]$ to X ($m \geq 1, j = 1, \dots, m$), extends to a map from $\Delta[m]$. Let us call $Kan(m, j)$ the Kan condition for the horn $\Lambda[m, j]$. A Kan simplicial set is therefore a simplicial set satisfying $Kan(m, j)$ for all $m \geq 1$ and $0 \leq j \leq m$. In the language of groupoids, the Kan condition corresponds to the possibility of composing various morphisms. For example, the existence of a composition for arrows is given by the condition $Kan(2, 1)$, whereas the composition of an arrow with the inverse of another is given by $Kan(2, 0)$ and $Kan(2, 2)$.



Note that the composition of two arrows is in general not unique, but any two of them can be joined by a 2-morphism h given by $Kan(3, 1)$.



Here, h ought to be a bigon, but since we don't have any bigons in a simplicial set, we view it as a triangle with one of its edges degenerate. The degenerate 1-simplex above z is denoted 1_z .

In an n -groupoid, the only well defined composition law is the one for n -morphisms. This motivates the following definition.

Definition 1.1. An n -groupoid ($n \in \mathbb{N} \cup \infty$) X is a simplicial set that satisfies $Kan(m, j)$ for all $0 \leq j \leq m \geq 1$ and $Kan!(m, j)$ for all $0 \leq j \leq m < n$, where


$Kan(m, j)$: Any map $\Lambda[m, j] \rightarrow X$ extends to a map $\Delta[m] \rightarrow X$.

$Kan!(m, j)$: Any map $\Lambda[m, j] \rightarrow X$ extends to a unique map $\Delta[m] \rightarrow X$.

An ∞ -groupoid will be called a homotopy type.

An n -group is an n -groupoid for which X_0 is a point. When $n = 2$, they are the same as the weak 2-group(oid)s in [Noo] but different from the various kinds of 2-group(oid)s or double groupoids in [BL04, BS76]². A usual groupoid (category with only isomorphisms) is equivalent to a 1-groupoid in the sense of Definition 1.1. Indeed, from a usual groupoid, one can form a simplicial set whose n -simplices are given by sequences of n composable arrows. This is a standard construction called the *nerve* of a groupoid and one can check that it satisfies the required Kan conditions.

On the other hand, a 1-groupoid X in the sense of Definition 1.1 gives us a usual groupoid with objects and arrows given respectively by the 0-simplices and 1-simplices of X . The unit is provided by the degeneracy $X_0 \rightarrow X_1$, the inverse and composition are given by the Kan conditions $Kan(2, 0)$, $Kan(2, 1)$ and $Kan(2, 2)$ as in (3), and the associativity is given by $Kan(3, 1)$ and $Kan!(2, 1)$.



² See [Hen] for an explanation of the relation with [BL04].

This motivates the corresponding definition in the differentiable category.

Definition 1.2. A Lie n -groupoid X ($n \in \mathbb{N} \cup \infty$) is a simplicial manifold that satisfies $Kan(m, j)$ for all $0 \leq j \leq m \geq 1$ and $Kan!(m, j)$ $0 \leq j \leq m < n$, where

$Kan(m, j)$: The restriction map $\text{hom}(\Delta[m], X) \rightarrow \text{hom}(\Lambda[m, j], X)$ is a surjective submersion.

$Kan!(m, j)$: The restriction map $\text{hom}(\Delta[m], X) \rightarrow \text{hom}(\Lambda[m, j], X)$ is a diffeomorphism.

A Lie ∞ -groupoid will be called a Lie homotopy type.

We view simplicial sets $\Delta[m]$ and $\Lambda[m, j]$ as simplicial manifolds with their discrete topology. Then $\text{hom}(S, X)$ denotes the set of homomorphisms of simplicial manifolds, with its natural topology, so $\text{hom}(\Delta[m], X)$ is just another name for X_m . However the fact that $\text{hom}(\Lambda[m, j], X)$ is a manifold is not obvious (see Section 2).

On the other hand, a *stacky Lie (SLie) groupoid* $\mathcal{G} \Rightarrow M$, following the concept of Weinstein (W-) groupoid in [TZ06a], is a groupoid object in the world of differentiable stacks with its base M an honest manifold (see Definition 3.1 for detailed axioms). When $\mathcal{G}$ is also a manifold, $\mathcal{G} \Rightarrow M$ is obviously a Lie groupoid. *W-groupoids*, which are *étale SLie* groupoids, provide a way to build the 1-1 correspondence with Lie algebroids.

Given these two higher generalizations of Lie groupoids, Lie n -groupoids and SLie groupoids, arising from different motivations and constructions, we ask the following questions:

- Are SLie groupoids the same as Lie n -groupoids for some n ?
- If not exactly, to which extent they are the same?
- Is there a way to also realize Lie n -groupoids as integration objects of Lie algebroids?

In this paper, we answer the two first questions by

Theorem 1.3. *There is a one-to-one correspondence between SLie (respectively W-) groupoids and Lie 2-groupoids (respectively Lie 2-groupoids whose X_2 is étale over $\text{hom}(\Lambda[2, j], X)$) modulo 1-Morita equivalences³ of Lie 2-groupoids.*

The last question will be answered positively in a future work [Zhu]:

Theorem 1.4. *Let A be a Lie algebroid and let $Lmor(-, -)$ be the space of Lie algebroid homomorphisms satisfying suitable boundary conditions. Then*

$$Lmor(T\Delta^2, A)/Lmor(T\Delta^3, A) \Rightarrow Lmor(T\Delta^1, A) \Rightarrow Lmor(T\Delta^0, A),$$

is a Lie 2-groupoid corresponding to the W-groupoid $\mathcal{G}(A)$ constructed in [TZ06a] under the correspondence in the above theorem.

For the first theorem, we (are forced to) develop Morita equivalence of Lie 2-groupoids, which is expected to be useful in the theory of 2-stacks and 2-gerbes and should correspond to Morita equivalence of SLie groupoids [BZ].

³Morita equivalences preserving X_0

We also find some technical improvements of the concept of SLie groupoids: it turns out that an SLie groupoid $\mathcal{G} \Rightarrow M$ always has a “good groupoid presentation” G of $\mathcal{G}$, which possesses a strict groupoid map $M \rightarrow G$. Moreover the condition on the inverse map could be simplified.

Acknowledgements: Here I would like to thank Henrique Bursztyn, Ezra Getzler, Alan Weinstein, for their hosts and very helpful discussions. I also thank Laurent Bartholdi and Marco Zambon for many editing suggestions. Finally I thank especially André Henriques who pointed out to me the potential correspondence of stacky groupoids and Lie 2-groupoids during the conference of “Groupoids and Stacks in Physics and Geometry” in CIRM-Luminy 2004. I owe a lot to discussions with André. He contributed the exact definitions of Lie n -groupoids and their equivalences, and also nice pictures!

2 Lie n -groupoids

In differential geometry, Lie groupoids have been studied a lot (See [CdSW99] for details). They are used to study foliations, and more recently orbifolds and differentiable stacks [Moe02] [BX]. Here we will try to convince the reader that it is fruitful to consider them within the context of Lie n -groupoids (Def. 1.2), especially if one wants to define and use sheaf cohomology. We shall also embed the category of manifolds in the category of Lie homotopy types by sending a manifold M to the constant simplicial manifold $\underline{M}$ with $\underline{M}_n = M$. Note that the constant simplicial sets (the ones whose only non-degenerate simplices are 0-dimensional) are exactly the 0-groupoids and they form a category which is equivalent to the category of sets. Similarly, the constant simplicial manifolds are exactly the Lie 0-groupoids. In the future, we shall abuse language and say “ X is a manifold” to mean “ X is a constant simplicial manifold”.

As promised after Def. 1.2, we first prove the important fact that $\text{hom}(\Lambda[j, m], X)$ is a manifold to make the definition complete. Since Lie n -groupoids are special Lie homotopy types (or you could call them Kan simplicial manifolds), we only have to show this fact for Lie homotopy types.

Lemma 2.1. *Let S be a finite k -dimensional collapsible simplicial set, namely one that can be obtained from the point by successively filling horns. Let X be a simplicial manifold that satisfies $\text{Kan}(m, j)$, $\forall 1 \leq m \leq k$, $j = 0, \dots, m$. Then the space $\text{hom}(S, X)$ is naturally a manifold.*

Here we assume that the spaces $\text{hom}(\Lambda[m, j], X)$ are known to be manifolds when $m \leq k$, for the condition $\text{Kan}(m, j)$ to make sense.

Proof. Let $S' \subset S$ be a collapsible sub-simplicial set such that S is obtained from S' by filling one horn. In other words, there is a push-out diagram

$$\begin{array}{ccc} S' & \xrightarrow{\quad} & S \\ \uparrow & \lrcorner & \uparrow \\ \Lambda[m, j] & \hookrightarrow & \Delta[m] \end{array} \quad ,$$

where the lower arrow is the inclusion. When applying $\text{hom}(_, X)$ to the above push-out, we get a pull-back diagram

$$\begin{array}{ccc} \text{hom}(S', X) & \longleftarrow & \text{hom}(S, X) \\ \downarrow & \lrcorner & \downarrow \\ \text{hom}(\Lambda[m, j], X) & \longleftarrow & \text{hom}(\Delta[m], X) = X_m \end{array}$$

By induction on the number of simplices of S we may assume that $\text{hom}(S', X)$ and $\text{hom}(\Lambda[m, j], X)$ are known to be manifolds. Moreover, by $Kan(m, j)$, the bottom arrow is a submersion. Therefore by transversality, $\text{hom}(S, X)$ is a manifold. $\square$

Since the horns $\Lambda[m, j]$ are collapsible, we have:

Corollary 2.2. *Let X be a Lie homotopy type, then $\text{hom}(\Lambda[m, j], X)$ is a manifold.*

2.1 The description of Lie 2-group(oid)s with finite data

Often the conventional way with only finite layers of data to understand Lie group(oid)s is more conceptual in differential geometry. Therefore it is worth giving also the finite description of Lie 2-groupoids. It turns out to be quite involved, however it provides a direct generalization of Lie groupoids and it is useful to pass to SLie groupoids. Moreover the version of 2-groupoid (not Lie) is completely analogous to it.

The equivalent description of Lie 2-groupoids uses only three layers $X_2 \rightrightarrows X_1 \rightrightarrows X_0$ and associative “3-multiplications” [Dus79]. Following the notion of simplicial manifolds, we call d_i^n and s_j^n the face and degeneracy maps between X_i ’s; they still satisfy the coherence condition in (1). To simplify the notation and match it with the definition of groupoids, we use the notation $\mathbf{t}$ for d_0^1 , $\mathbf{s}$ for d_1^1 and e for s_0^0 . Then we can safely omit the upper indices for d_i^2 ’s and s_i^1 ’s. Actually we will omit the upper indices whenever it does not cause confusion. Similarly to the horn spaces $\text{hom}(\Lambda[m, j], X)$, given only these three layers, we define $\Lambda(X)_{m,j}$ to be the space of m elements in X_{m-1} gluing along elements in X_{m-2} to a horn shape without the j -th face.

$$\begin{array}{ccccccc} \bullet & \bullet & \begin{array}{c} \bullet \\ \nearrow \searrow \\ \bullet \end{array} & \begin{array}{c} \bullet \\ \nearrow \searrow \\ \bullet \end{array} & \begin{array}{c} \bullet \\ \nearrow \searrow \\ \bullet \end{array} & \begin{array}{c} \bullet \quad \bullet \\ \nearrow \quad \searrow \\ \bullet \quad \bullet \end{array} & \begin{array}{c} \bullet \quad \bullet \\ \nearrow \quad \searrow \\ \bullet \quad \bullet \end{array} \\ \Lambda(X)_{1,1} & \Lambda(X)_{1,0} & \Lambda(X)_{2,2} & \Lambda(X)_{2,1} & \Lambda(X)_{2,0} & \Lambda(X)_{3,3} & \Lambda(X)_{3,2} \dots \end{array}$$

Here one imagines each j -dimensional face as an element in X_j . For example,

$$\begin{aligned} \Lambda(X)_{2,2} &= X_1 \times_{\mathbf{s}, X_0, \mathbf{s}} X_1, \quad \Lambda(X)_{2,1} = X_1 \times_{\mathbf{t}, X_0, \mathbf{s}} X_1, \quad \Lambda(X)_{2,0} = X_1 \times_{\mathbf{t}, X_0, \mathbf{t}} X_1, \\ &\dots, \quad \Lambda(X)_{3,0} = (X_2 \times_{d_2, X_1, d_1} X_2) \times_{d_1 \times d_2, \Lambda(X)_{2,0}, d_1 \times d_2} X_2. \end{aligned}$$

We remark that item (1a) and (1b) in the proposition-definition below imply that $\Lambda(X)_{2,j}$ ’s and $\Lambda(X)_{3,j}$ ’s are manifolds. Then with this condition we can define *3-multiplications*

as smooth maps $m_i : \Lambda(X)_{3,i} \rightarrow X_2$, $i = 0, \dots, 3$. With 3-multiplications, there are natural maps between $\Lambda(X)_{3,j}$'s. For example,

$$\Lambda(X)_{3,0} \rightarrow \Lambda(X)_{3,1}, \quad \text{by } (\eta_1, \eta_2, \eta_3) \rightarrow (m_0(\eta_1, \eta_2, \eta_3), \eta_2, \eta_3).$$

It is reasonable to ask them to be isomorphisms. In fact, set theoretically, this simply says that the following four equations are equivalent to each other:

$$\begin{aligned} \eta_0 &= m_0(\eta_1, \eta_2, \eta_3), & \eta_1 &= m_1(\eta_0, \eta_2, \eta_3), \\ \eta_2 &= m_2(\eta_0, \eta_1, \eta_3), & \eta_3 &= m_3(\eta_0, \eta_1, \eta_2). \end{aligned}$$

Proposition-Definition 2.3. A Lie 2-groupoid can be also described by three layers $X_2 \rightrightarrows X_1 \rightrightarrows X_0$ and the following data:

1. the face and degeneracy maps d_i^n and s_i^n satisfying (1) for $n = 1, 2$ as explained above, and
 - (a) [1-Kan] $\mathbf{t}$ and $\mathbf{s}$ are surjective submersions;
 - (b) [2-Kan] $d_0 \times d_2 : X_2 \rightarrow \Lambda(X)_{2,1} = X_1 \times_{\mathbf{t}, X_0, \mathbf{s}} X_1$, $d_0 \times d_1 : X_2 \rightarrow \Lambda(X)_{2,2} = X_1 \times_{\mathbf{s}, X_0, \mathbf{s}} X_1$, and $d_1 \times d_2 : X_2 \rightarrow \Lambda(X)_{2,0} = X_1 \times_{\mathbf{t}, X_0, \mathbf{t}} X_1$ are surjective submersions.
2. smooth maps (3-multiplications),

$$m_i : \Lambda(X)_{3,i} \rightarrow X_2 \quad i = 0, \dots, 3.$$

such that

- (a) the induced morphisms (by m_j as above) $\Lambda(X)_{3,i} \rightarrow \Lambda(X)_{3,j}$ are all isomorphisms;
- (b) m_i 's are compatible with the face and degeneracy maps:

$$\begin{aligned} \eta &= m_1(\eta, s_0 \circ d_1(\eta), s_0 \circ d_2(\eta)) \text{ (which is equivalent to } \eta = m_0(\eta, s_0 \circ d_1(\eta), s_0 \circ d_2(\eta))), \\ \eta &= m_2(s_0 \circ d_0(\eta), \eta, s_1 \circ d_2(\eta)) \text{ (which is equivalent to } \eta = m_1(s_0 \circ d_0(\eta), \eta, s_1 \circ d_2(\eta))), \\ \eta &= m_3(s_1 \circ d_0(\eta), s_1 \circ d_1(\eta), \eta) \text{ (which is equivalent to } \eta = m_2(s_1 \circ d_0(\eta), s_1 \circ d_1(\eta), \eta)). \end{aligned} \tag{5}$$

- (c) m_i 's are associative, that is for a 5-simplex $(0, 1, 2, 3, 4)$,

$$\begin{array}{c} \begin{array}{ccccc} & & 0 & & \\ & \nearrow & \uparrow & \nwarrow & \\ & 4 & \xrightarrow{\quad} & 1 & \\ & \nwarrow & \searrow & \nearrow & \\ 3 & & & & 2 \end{array} \end{array} \tag{6}$$

if we are given faces $(0, i, 4)$'s and $(0, i, j)$'s in X_2 , where $i, j \in \{1, 2, 3\}$, then the following two methods to determine the face $(1, 2, 3)$ give the same element in X_2 :

- i. $(1, 2, 3) = m_0((0, 2, 3), (0, 1, 3), (0, 1, 2));$
- ii. we first obtain $(i, j, 4)$ using m_i 's on $(0, i, 4)$'s, then we have

$$(1, 2, 3) = m_3((2, 3, 4), (1, 3, 4), (1, 2, 4)).$$

Remark 2.4. Here we notice that the surjectivity of the maps in the 2-Kan condition (1b) insure the existence of a usual (2-) multiplication $m : X_1 \times_{\mathbf{t}, X_0, \mathbf{s}} X_1$ and an inverse $i : X_1 \rightarrow X_1$ as explained in the introduction. For example since $d_0 \times d_2 : X_2 \rightarrow \Lambda(X)_{2,1}$ is surjective we could take a section of it and compose it with $d_1 : X_2 \rightarrow X_1$. This gives a multiplication map m , which is furthermore smooth if we take a smooth section. This can always be achieved by passing to an equivalent Lie 2-groupoid (see Section 6.1). However it is usually not associative on the nose, but only up to elements in X_2 . One could make it strictly associative by passing to an infinite dimensional model. But for us the important example coming from integrating Lie algebroids does not satisfy associativity on the nose. As stated in Theorem 1.4, the Lie 2-groupoid has $X_1 = Lmor(T\Delta^1, A)$, which is the space of A -paths satisfying the boundary condition $a(0) = a(1) = 0$. The 2-multiplication m is simply the concatenation of A -paths which is only associative up to A -homotopies. Moreover we can not simply mod out A -homotopies from X_1 since the resulting space might not be a smooth manifold anymore.

On the other hand, only having a usual 2-multiplication m and an inverse map i , it is not guaranteed that the maps in the 2-Kan condition (1b) are submersions even when m and i are smooth. But being submersions is in turn very important to prove that $X_{n \geq 3}$ are smooth manifolds. Hence in the differentiable category, we can not replace the 2-Kan condition by a usual 2-multiplication and an inverse.

The nerve of $X_2 \rightrightarrows X_1 \rightrightarrows X_0$

To show that what we defined just now is the same as Def. 1.2, we form the *nerve* of a Lie 2-groupoid $X_2 \rightrightarrows X_1 \rightrightarrows X_0$ in Prop-Def. 2.3. We first define

$$X_3 = \{(\eta_0, \eta_1, \eta_2, \eta_3) : \eta_0 = m_0(\eta_1, \eta_2, \eta_3), (\eta_1, \eta_2, \eta_3) \in \Lambda(X)_{3,0}\}.$$

Then $X_3 \cong \Lambda(X)_{3,0}$ is a manifold. Moreover, we have the obvious face and degeneracy maps between X_3 and X_2 ,

$$\begin{aligned} d_i^3(\eta_0, \eta_1, \eta_2, \eta_3) &= \eta_i, \quad i = 0, \dots, 3 \\ s_0^2(\eta) &= (\eta, \eta, s_0 \circ d_1(\eta), s_0 \circ d_2(\eta)), \\ s_1^2(\eta) &= (s_0 \circ d_0(\eta), \eta, \eta, s_1 \circ d_2(\eta)), \\ s_2^2(\eta) &= (s_1 \circ d_0(\eta), s_1 \circ d_1(\eta), \eta, \eta). \end{aligned}$$

The coherency (5) insures that $s_i^2(\eta) \in X_3$. It is also not hard to see that these maps together with $d_i^{\leq 2}$'s and $s_i^{\leq 1}$'s satisfy (1) for $n \leq 3$.

Then the nerve can be easily described as the simplicial manifold

$$X = \text{cosk}_2(\text{sk}_2(X_3 \rightarrow X_2 \rightarrow X_1 \rightarrow X_0)).$$

More concretely, X_n is made up by those n -simplices whose 2-faces are elements of X_2 and such that each set of four 2-faces gluing together as a 3-simplex is an element of X_3 . That is

$$X_n = \{f \in \text{hom}_2(sk_2(\Delta_n), X_2) | f \circ (d_0 \times d_1 \times d_2 \times d_3)(sk_3(\Delta_n)) \subset X_3\},$$

where hom_2 denotes the homomorphisms restricted to the 0,1,2 level and X_2 is understood as the tower $X_2 \Rightarrow X_1 \Rightarrow X_0$ with all degeneracy and face maps. Then there are obvious face and degeneracy maps which naturally satisfy (1).

However what is nontrivial and special in the differentiable category is that the associativity of m_i 's assures that X_n is a manifold. We prove this by an inductive argument. Let $S_j[n] = sk_2(\Lambda[n, j])$. It is the contractible simplicial set whose sub-faces all contain the vertex j and whose only non-degenerate faces are of dimension 0, 1 and 2. Then similarly to Lemma 2.1, we show that $\text{hom}_2(S_j[n], X_2)$ is a manifold. Since $S_j[n]$ is constructed by adding 0,1,2 dimensional faces, it is formed by the procedure

$$\begin{array}{ccc} S' & \longrightarrow & S \\ \uparrow & & \uparrow \\ \Lambda[n, j] & \longrightarrow & \Delta[n] \end{array}$$

with $n \leq 2$. The dual pull-back diagram shows that $\text{hom}_2(S_j[n], X_2)$ is a manifold by induction

$$\begin{array}{ccc} \text{hom}_2(S', X_2) & \longleftarrow & \text{hom}_2(S, X_2) \\ \downarrow & & \downarrow \\ \text{hom}_2(\Lambda[n, j], X_2) & \longleftarrow & \text{hom}_2(\Delta[n], X_2), \end{array}$$

since $\text{hom}_2(\Delta[n], X_2) \rightarrow \text{hom}_2(\Lambda[n, j], X_2)$ are surjective submersions by item (1a) and (1b) in the last Prop-Def.

Next we use induction to show that $X_n = \text{hom}_2(S_0[n], X_2)$. Similarly we will have $X_n = \text{hom}_2(S_j[n], X_2)$. It is clear that $\tilde{f} \in X_n$ restricts to $\tilde{f}|_{S_0[n]} \in \text{hom}_2(S_0[n], X_2)$. We only have to show that $f \in \text{hom}_2(S_0[n], X_2)$ extends uniquely to $\tilde{f} \in X_n$. It is certainly true for $n = 0, 1, 2, 3$ just by definition. Suppose $X_{n-1} = \text{hom}_2(S_0[n-1], X_2)$. Then to get $f \in \text{hom}_2(S_0[n], X_2)$ from $f' \in \text{hom}_2(S_0[n-1], X_2)$, we add a new point n and $n-1$ new faces $(0, i, n)$, $i \in \{1, 2, \dots, (n-1)\}$ and dye them red⁴. Using 3-multiplication m_0 , we can determine face (i, j, n) by $(0, i, n)$, $(0, j, n)$ and $(0, i, j)$ and dye these newly decided faces blue. Now we want to see that each four faces attached together are in X_3 , then f is extended to $\tilde{f} \in X_n$. We consider various cases:

1. if none of the four faces contains the vertex n , then by the induction condition, they are in X_3 .
2. if one of the four faces contains n , then there are three faces containing n , we again have two sub-cases:
 - (a) if those three faces contain only one blue face of the form (i, j, n) , $i, j \in \{1, \dots, (n-1)\}$, then the four faces must contain three red faces and one blue face. According to our construction, these four faces are in X_3 ;

⁴More precisely, they are the image of these under the map f .

- (b) if those three faces contains more than one blue face, then they must contain exactly three blue faces. Then according to associativity (inside the 5-gon $(0, i, j, k, n)$), these four faces are also in X_3 .

Now we finish the induction, hence X_n is a manifold and it is determined by the first three layers. So we have $\text{hom}(\Lambda[n, j], X) = \text{hom}_2(sk_2(\Lambda[n, j]), X_2) = X_n$ and,

Proposition 2.5. *The nerve X of a Lie 2-groupoid $X_2 \rightrightarrows X_1 \rightrightarrows X_0$ as in Prop-Def. 2.3 is a Lie 2-groupoid as in Def. 1.2.*

Proposition 2.6. *The first three layers of a Lie 2-groupoid as in Definition 1.2 is a Lie 2-groupoid as in Prop-Def. 2.3.*

Proof. The proof is more complicated and similar to the case of 1-groupoids in the introduction. Here we point out that the 3-multiplications m_j are given by $Kan(3, j)$ and the associativity is given by $Kan!(3, 0)$ and $Kan(4, 0)$. $\square$

3 SLie groupoids

First we give the precise definition of SLie groupoids and W-groupoids. This generalizes and completes the notion of Weinstein (W-) groupoids in [TZ06a]. For example, we add some new axioms on the level of 2-morphisms, and on the other hand, find that some other axioms could be replaced or simplified. Then we point out some direct implications from the definition. The notion of stacks has been extensively studied in algebraic geometry for the past few decades. However stacks can also be defined over other categories, such as the category of topological spaces and category of smooth manifolds (see for example [AGV72] [Pro96] [Vis02] [BX] [Met]). We refer the readers to the latter two references for the concepts we use here, such as (étale) differentiable stacks, their fibre products, immersion maps between them. But for *surjective submersion*, we adopt the definition in [TZ06a] which is a bit different. $f : \mathcal{X} \rightarrow \mathcal{Y}$ is a submersion if $X \times_{\mathcal{Y}} Y \rightarrow Y$ is a submersion where X and Y are charts of $\mathcal{X}$ and $\mathcal{Y}$ respectively. f is further a surjective submersion if it is an epimorphism of differentiable stacks.

Definition 3.1 (SLie (respectively W-) groupoid). A stacky Lie or SLie (respectively Weinstein or W-) groupoid over a manifold M consists of the following data:

1. a differentiable (respectively an étale differentiable) stack $\mathcal{G}$;
2. (source and target) maps $\bar{s}, \bar{t} : \mathcal{G} \rightarrow M$ which are surjective submersions between differentiable stacks;
3. (multiplication) a map m : the fibre product $\mathcal{G} \times_{\bar{s}, M, \bar{t}} \mathcal{G} \rightarrow \mathcal{G}$, satisfying the following properties:
 - (a) $\bar{t} \circ m = \bar{t} \circ pr_1$, $\bar{s} \circ m = \bar{s} \circ pr_2$, where $pr_i : \mathcal{G} \times_{\bar{s}, M, \bar{t}} \mathcal{G} \rightarrow \mathcal{G}$ is the i -th projection $\mathcal{G} \times_{\bar{s}, M, \bar{t}} \mathcal{G} \rightarrow \mathcal{G}$;
 - (b) associativity up to a 2-morphism, i.e. there is a 2-morphism a between maps $m \circ (m \times id)$ and $m \circ (id \times m)$;

- (c) the 2-morphism a satisfies a higher coherence described as following: let the 2-morphisms on the each face of the cubes be a_i ⁵ arranged in the following way: front face (the one with the most $\mathcal{G}$'s) a_1 , back a_5 ; up a_4 , down a_2 ; left a_6 , right a_3 ,

$$\begin{array}{ccccc}
 & & \mathcal{G} \times \mathcal{G} \times \mathcal{G} & & \\
 & & M & & \\
 & \nearrow^{id \times id \times m} & & \searrow_{m \times id} & \\
 \mathcal{G} \times \mathcal{G} \times \mathcal{G} \times \mathcal{G} & & \mathcal{G} \times \mathcal{G} \times \mathcal{G} & & \mathcal{G} \times \mathcal{G} \\
 M & M & M & M & M \\
 & \searrow_{m \times id \times id} & \nearrow^{id \times m} & \searrow_{id \times m} & \\
 & \mathcal{G} \times \mathcal{G} \times \mathcal{G} & & & \\
 & M & & & \\
 \downarrow^{id \times m \times id} & & \downarrow^{id \times m} & & \downarrow^m \\
 \mathcal{G} \times \mathcal{G} \times \mathcal{G} & & \mathcal{G} \times \mathcal{G} & & \mathcal{G} \\
 M & M & M & & \\
 & \nearrow^{id \times m} & \searrow_{m \times id} & & \\
 & \mathcal{G} \times \mathcal{G} & & & \\
 & M & & & \\
 & \searrow_{m \times id} & \nearrow^m & & \\
 & \mathcal{G} \times \mathcal{G} & & & \\
 & M & & &
 \end{array}$$

We require

$$(a_6 \times id) \circ (id \times a_2) \circ (a_1 \times id) = (id \times a_5) \circ (a_4 \times id) \circ (id \times a_3).$$

4. (identity section) a morphism (respectively an immersion) $\bar{e}: M \rightarrow \mathcal{G}$ such that
- (a) the following identities

$$m \circ ((\bar{e} \circ \bar{\mathbf{t}}) \times id) = id, \quad m \circ (id \times (\bar{e} \circ \bar{\mathbf{s}})) = id,$$

hold⁶ up to 2-morphisms b_l and b_r . Or equivalently there are two 2-morphisms

$$\begin{array}{ccc}
 m \circ (id \times \bar{e}) \xrightarrow{b_r} pr_1 : \mathcal{G} \times_{\bar{\mathbf{s}}, M} M \rightarrow \mathcal{G}, & m \circ (\bar{e} \times id) \xrightarrow{b_l} pr_2 : M \times_{M, \bar{\mathbf{t}}} \mathcal{G} \rightarrow \mathcal{G}, \\
 g\bar{e}(y) \rightarrow g & \bar{e}(x)g \rightarrow g
 \end{array}$$

where $y = \bar{\mathbf{s}}(g)$ and $x = \bar{\mathbf{t}}(g)$.

- (b) We require the composed 2-morphism below, with $y = \bar{\mathbf{s}}(g_2)$,

$$g_1 g_2 \xrightarrow{b_r^{-1}} (g_1 g_2) \bar{e}(y) \xrightarrow{a} g_1 (g_2 \bar{e}(y)) \xrightarrow{b_r} g_1 g_2$$

to be the identity.⁷

⁵All the a_i 's are generated by a , except that a_4 is id .

⁶In particular, by combining with the surjectivity of $\bar{\mathbf{s}}$ and $\bar{\mathbf{t}}$, one has $\bar{\mathbf{s}} \circ \bar{e} = id$, $\bar{\mathbf{t}} \circ \bar{e} = id$ on M . In fact if $x = \bar{\mathbf{t}}(g)$, then $\bar{e}(x) \cdot g = g$ and $\bar{\mathbf{t}} \circ m = \bar{\mathbf{t}} \circ pr_1$ imply that $\bar{\mathbf{t}}(\bar{e}(x)) = \bar{\mathbf{t}}(g) = x$.

⁷Forming this in the language of differentiable geometry, we notice that $pr_1 \circ (m \times id)$ and $m \circ (pr_1 \times pr_2)$ are the same map from $\mathcal{G} \times_M \mathcal{G} \times_M M$ to $\mathcal{G}$, but as the diagram indicates,

$$\begin{array}{ccc}
 \mathcal{G} \times_M \mathcal{G} \times_M M & \xrightarrow{m \times id} & \mathcal{G} \times_M M \\
 \downarrow^{id \times (m \circ (id \times \bar{e}))} \downarrow_{pr_1 \times pr_2} & & \downarrow_{pr_1} \downarrow_{m \circ (id \times \bar{e})} \\
 \mathcal{G} \times_M \mathcal{G} & \xrightarrow{m} & \mathcal{G},
 \end{array} \tag{7}$$

(c) similarly with $x = \bar{\mathbf{t}}(g_1)$,

$$g_1 g_2 \xrightarrow{b_l^{-1}} \bar{e}(x)(g_1 g_2) \xrightarrow{a^{-1}} (\bar{e}(x)g_1)g_2 \xrightarrow{b_l} g_1 g_2$$

is the identity;

(d) with $y = \bar{\mathbf{s}}(g)$ and $x = \bar{\mathbf{t}}(g)$,

$$g \xrightarrow{b_l^{-1}} \bar{e}(x)g \xrightarrow{b_r^{-1}} (\bar{e}(x)g)\bar{e}(y) \xrightarrow{b_l} g\bar{e}(y) \xrightarrow{b_r} g,$$

is the identity.

5. (inverse) an isomorphism of differentiable stacks $\bar{i}: \mathcal{G} \rightarrow \mathcal{G}$ such that, up to 2-morphisms, the following identities

$$m \circ (\bar{i} \times id \circ \Delta) = \bar{e} \circ \bar{\mathbf{s}}, \quad m \circ (id \times \bar{i} \circ \Delta) = \bar{e} \circ \bar{\mathbf{t}},$$

hold, where Δ is the diagonal map: $\mathcal{G} \rightarrow \mathcal{G} \times \mathcal{G}$.

Remark 3.2. A W-groupoid is simply an étale SLie groupoid. This definition of W-groupoid is different from the one in [TZ06a] in two aspects: one is that here we add various higher coherences on 2-morphisms which make the definition more restricting but still allow the W-groupoids $\mathcal{G}(A)$ and $\mathcal{H}(A)$, which are the integration objects of the Lie algebroid A constructed in [TZ06a]; for the other see Remark 3.6. On the other hand, we do not add higher coherences for the 2-morphisms involving the inverse map. This is because the inverse map can be removed from the definition. See Section 8.

Moreover, we notice that the 2-morphisms $\bar{e}(x) \cdot \bar{e}(x) \xrightarrow[b_r]{b_l} \bar{e}(x)$ are the same because they are basically 2-morphisms between morphisms on a manifold M . With some patience, we can check that the list of coherences on 2-morphisms given here generates all the possible coherences on these 2-morphisms. We also notice that the cube condition (3c) is the differential version of the pentagon condition

$$[((gh)k)l \rightarrow (g(hk))l \rightarrow g((hk)l) \rightarrow g(h(kl))] = [((gh)k)l \rightarrow (gh)(kl) \rightarrow g(h(kl))].$$

3.1 Good charts

Given an SLie groupoid $\mathcal{G} \Rightarrow M$, the identity map $\bar{e}: M \rightarrow \mathcal{G}$ corresponds to an Hilsum-Skandalis (H.S.) morphism from $M \Rightarrow M$ to $G_1 \Rightarrow G_0$ for some presentation of $\mathcal{G}$. But it is not clear whether M embeds into G_0 . It is not even obvious whether there is a map $M \rightarrow G_0$. In general, one could ask: if there is a map from a manifold M to a differentiable

they are related also via a sequence of 2-morphisms:

$$pr_1 \circ (m \times id) \xrightarrow{b_r^{-1} \odot id} m \circ (id \times \bar{e}) \circ (m \times id) \xrightarrow{a} m \circ (id \times (m \circ (id \times \bar{e}))) \xrightarrow{id \odot (id \times b_r)} m \circ (pr_1 \times pr_2) \quad (8)$$

We require the composed 2-morphisms be id , that is

$$(id \odot (id \times b_r)) \circ a \circ (b_r^{-1} \odot id) = id.$$

Because of uniqueness and $g_{ij} \cdot g_{jk} = g_{ik}$, we have $\bar{g}_{ij} \cdot \bar{g}_{jk} = \bar{g}_{ik}$ on the open subsets $\bar{U}_{ijk} := \{(\bar{g}_{ij}, \bar{g}_{jk}, \bar{g}_{ik}) : \text{when } \bar{g}_{ij} \cdot \bar{g}_{jk} \text{ exists and in } \bar{U}_{ik}\}$. Then

$$e_i(V_i \cap V_j \cap V_k) \subset U_{ijk} := \mathbf{t}(\text{Im}(\bar{U}_{ijk} \rightarrow \bar{U}_{ij})) \subset U_{ij} \cap U_{ik} \subset U_i,$$

and similarly for j and k . Therefore with these U 's we are in the situation of a *germ of manifolds* of M defined as below.

A *germ of manifolds at a point m* is a series of manifolds U_i 's containing m such that U_i agrees with U_j in a smaller open set $(m \in) U_{ij} \subset U_i$ by $x \sim f_{ij}(x)$. A *compatible riemannian metric* of a germ of manifolds consists of a riemannian metric g^i on each U_i such that two such riemannian metrics g^i and g^j on U_i and U_j agree with each other in the sense that $g^i(x) = g^j(f_{ij}(x))$ in a smaller open set (possibly a subset of U_{ij}). With this one can define the exponential map $\exp$ at m using the usual exponential map of a riemannian manifold provided the germ is finite, namely there are finite many manifolds in the germ (which is true in our case, since V_i intersects finite other V_j 's). Then $\exp$ gives a Hausdorff manifold containing m .

If a series of locally finite manifolds U_i 's and morphisms f_{ij} 's form a germ of manifolds for every point of a manifold M , we call it a *germ of manifolds of M* . Here local finiteness means that any open set in M is contained in finitely many U_i 's and M has the topology induced by the U_i 's, that is $M \cap U_i$ is open in M . We can always endow each of them with a compatible riemannian metric, beginning with any riemannian metric g^i on U_i and modifying it to the sum $g^i(x) := \sum_{k, x \in U_{ik}} g^k(f_{ik}(x))$ (with $f_{ii}(x) = x$) at each point $x \in U_i$. In this situation, one can take a tubular neighborhood U of M by the $\exp$ map of the germ. Then U is a Hausdorff manifold.

Applying the above construction to our situation, we have a Hausdorff manifold $U \supset M$ with the same dimension as G_0 . U is basically glued by small enough open subsets $\tilde{U}_i = U \cap U_i$ containing V_i 's along $\tilde{U}_{ij} := U \cap U_{ij}$ so that the gluing result U is still a Hausdorff manifold. Therefore U is presented by $\sqcup \tilde{U}_{ij} \Rightarrow \sqcup \tilde{U}_i$ which maps to $G_1 \Rightarrow G_0$ via $U_{ij} \cong \bar{U}_{ij} \hookrightarrow G_1$. So there is a map $\pi : U \rightarrow \mathcal{G}$. Since $\tilde{U}_i \rightarrow G_0$ are étale maps, by the technical lemma below, π is a representable étale map. $\square$

Lemma 3.4. *Given a manifold X and an (étale) differentiable stack $\mathcal{Y}$, a map $f : X \rightarrow \mathcal{Y}$ is an (étale) representable submersion if and only if there exists an (étale) chart Y_0 of $\mathcal{Y}$ such that the induced local maps $X_i \rightarrow Y_0$ are (étale) submersions, where $\{X_i\}$ is an open covering of X .*

Proof. For any $V \rightarrow \mathcal{Y}$, $X_i \times_{\mathcal{Y}} V = X_i \times_{Y_0} Y_0 \times_{\mathcal{Y}} V$ is representable and $X_i \times_{\mathcal{Y}} V \rightarrow V$ is an (étale) submersion since $X_i \rightarrow Y_0$ and $Y_0 \rightarrow \mathcal{Y}$ are representable (étale) submersions. Since X_i glue together to X , $X_i \times_{\mathcal{Y}} V$ with the inherited gluing maps glue to a manifold $X \times_{\mathcal{Y}} V$. Since being an (étale) submersion is a local property, $X \times_{\mathcal{Y}} V \rightarrow V$ is an (étale) submersion.

$$\begin{array}{ccc}
 X_i \times_{\mathcal{Y}} V & & X_i \\
 \downarrow & \searrow & \downarrow \\
 & X_k \times_{\mathcal{Y}} V & X_k \\
 & \swarrow & \swarrow \\
 X_j \times_{\mathcal{Y}} V & & X_j
 \end{array}
 \quad \Leftarrow
 \quad
 \begin{array}{ccc}
 X_i & & X_i \\
 \downarrow & \searrow & \downarrow \\
 & X_k & X_k \\
 & \swarrow & \swarrow \\
 X_j & & X_j
 \end{array}$$

□

Remark 3.5. If $\bar{e}$ is the identity map of W-groupoid $\mathcal{G} \Rightarrow M$, then an open neighborhood of M in U has an induced local groupoid structure from the stacky groupoid structure [TZ06a, Section 5].

Remark 3.6. The following assumption in the original definition of W-groupoids becomes obviously unnecessary after Lemma 3.3:

“Moreover, restricting to the identity section, the above 2-morphisms between maps are the *id* 2-morphisms. Namely, for example, the 2-morphism α induces the *id* 2-morphism between the following two maps:

$$m \circ ((m \circ (\bar{e} \times \bar{e} \circ \delta)) \times \bar{e} \circ \delta) = m \circ (\bar{e} \times (m \circ (\bar{e} \times \bar{e} \circ \delta)) \circ \delta),$$

where δ is the diagonal map: $M \rightarrow M \times M$.”

This is unnecessary because $(M \Rightarrow M)$ embeds into $(G_1 \Rightarrow G_0)$ on both levels of the groupoid via $\bar{e}$ and the restricted maps become maps between manifolds and therefore there are no nontrivial 2-morphisms between them.

We further prove the same lemma in the non-étale case.

Lemma 3.7. *For a morphism $\bar{e} : M \rightarrow \mathcal{G}$ from a manifold M to a differentiable stack $\mathcal{G}$, there is a chart G_0 of $\mathcal{G}$ such that $\bar{e}$ lifts to an embedding $e : M \rightarrow G_0$. We call also such groupoid presentations good presentations.*

Proof. We follow the proof of the étale case, but replace “étale” with “submersion”. We need a U with a representable submersion to $\mathcal{G}$ and an embedding of M into U . There are two differences: first, V_i embeds in V'_i instead of being isomorphic to it, and we do not have an embedding $V'_i \hookrightarrow G_0$; second, since $G_1 \Rightarrow G_0$ is not étale, the bisection g_{ij} does not extend uniquely to some $\bar{g}_{ij}$ and we can not have the cocycle condition immediately.

The first difference is easy to compensate: given any morphism $f : N_1 \rightarrow N_2$, we can always view it as a composition of an embedding and a submersion as $N_1 \xrightarrow{id \times f} N_1 \times N_2 \xrightarrow{pr_2} N_2$. In our case, we have the following decomposition $M \times_{\mathcal{G}} G_0 \hookrightarrow H_0 \twoheadrightarrow G_0$, then we use the pull-back groupoid $H_1 := G_1 \times_{G_0 \times G_0} H_0 \times H_0$ over H_0 to replace G . Thus we obtain an embedding of $V'_i \rightarrow H_0$ and thus an embedding $V_i \rightarrow H_0$. Then since $H_1 \Rightarrow H_0$ is Morita equivalent to $G_1 \Rightarrow G_0$, we just have to replace G by H or call H our new G . It was not possible to do so in the étale case since H_0 might not be an étale chart of $\mathcal{G}$.

For the second difference, first of all we could assume M to be connected to construct such U . Otherwise we take the disjoint union of U ’s of each connected component of M .

Then take any V_i and consider all the charts V_j ’s intersecting V_i . We choose $\bar{g}_{ij}$ extending g_{ij} on an open set $\bar{U}_{ij}$. As before we define the open sets U_i , U_j ’s, and U_{ij} . Then for V_j and $V_{j'}$ both intersecting V_i , we choose $\bar{g}_{jj'}$ to be the one extending (see below) $\bar{g}_{ij}^{-1} \bar{g}_{ij'}$ with $\mathbf{s}(\bar{g}_{ij}^{-1} \bar{g}_{ij'})$ in the triple intersection $\bar{g}_{ij'}^{-1} \cdot (\bar{g}_{ij} \cdot U_j) \cap U_{j'}$ where multiplication applies when it can. Since $\bar{g}$ ’s are local bisections, $\bar{g} \cdot$ is an isomorphism. Identifying via these isomorphisms, we view and denote the above intersection as $U_{j'ij}$ for simplicity.

Now we clarify in which sense and why the extension always exists. Let us assume $\dim M = m$, $\dim G_i = n_i$. Here we identify V_j with its embedded image in G_0 . Then since we are dealing with local charts, we might assume that both $\mathbf{t}$ and $\mathbf{s}$ of $G_1 \Rightarrow G_0$ are just projections from $\mathbb{R}^{n_1}$ to $\mathbb{R}^{n_0}$. A section of $\mathbf{s}$ is a vector valued function $\mathbb{R}^{n_0} \rightarrow \mathbb{R}^{n_1}/\mathbb{R}^{n_0}$, and it being a bisection, namely also a section of $\mathbf{t}$, is an open condition. We can always perturb a section to get a bisection. If we can extend $\bar{g}_{ij}^{-1} \bar{g}_{ij'}$ and $g_{jj'}$ from $U_{j'ij} \cup e_{j'}(V_j \cap V_{j'})$ to a

bisection $\bar{g}_{jj'}$ such that $\{\mathbf{s}(\bar{g}_{jj'})\}$ is an open set in $U_{j'}$, then we obtain a bisection $\bar{g}_{jj'}$ from $U_{j'j} := \bar{g}_{jj'}^{-1}(\{\mathbf{t}(\bar{g}_{jj'})\} \cap U_j)$ to $U_{jj'} := \{\mathbf{t}(\bar{g}_{jj'})\} \cap U_j$. It is easy to see that $U_{jj'} \cong U_{j'j}$ are open in G_0 since $\{\mathbf{t}(\bar{g}_{jj'})\} \cong \{\mathbf{s}(\bar{g}_{jj'})\}$.

Therefore we are done as long as we can extend a smooth function f from the union of an open submanifold O with a closed submanifold V of an open set $B \subset \mathbb{R}^{n_0}$ to the whole B . Since V is closed, using its tubular neighborhood and partition of unity, we can first extend f from V to B as $\tilde{f}$. Then $f_1 = f - \tilde{f}|_{O \cup V}$ is 0 on V . We shrink the open set O a little bit to O_i such that $V \cap O \subset O_2 \subset O_1 \subset O$. Then we always have a smooth function p on B with $p|_{\bar{O}_2} = 1$ and $p|_{B-O_1} = 0$. Then the extension function $\tilde{f}_1$ is defined by

$$\tilde{f}_1(x) = \begin{cases} f_1(x) \cdot p(x) & x \in O, \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that $\tilde{f}_1$ is smooth and it agrees with f_1 on O_2 and V because $V - O_2 = V - O_1 \subset B - O_1$ and $p|_{V-O_2} = 0$. Hence $\tilde{f} + \tilde{f}_1$ extends $f|_{O_2 \cup V}$. Now we extend the $\bar{g}_{ij}^{-1}\bar{g}_{ij'}$'s to $\bar{g}_{jj'}$'s, then the $\bar{g}$'s satisfy the cocycle condition on smaller open sets of the triple intersections $U_{j'ij}$ by construction.

Then we view $V_i \cup (\cup_{j: V_i \cap V_j \neq \emptyset} V_j)$ as one chart. Notice that a connected manifold is path connected. Also notice that we didn't use any topological property of V_i or U_i . This construction will eventually extend to the whole manifold M and obtain the desired $\bar{g}_{ij}$'s. Therefore we are again in the situation of a germ of manifolds and we can apply the proof of Lemma 3.3 to get the result. $\square$

4 From SLie-groupoids to Lie 2-groupoids

Suppose $\mathcal{G} \Rightarrow M$ is an SLie groupoid, in this section we construct a Lie 2-groupoid $X_2 \Rightarrow X_1 \Rightarrow X_0$ corresponding to it. When $\mathcal{G} \Rightarrow M$ is a W-groupoid, the corresponding Lie 2-groupoid is 2-étale, that is the maps $X_2 \rightarrow \text{hom}(\Lambda[2, j], X)$ are étale for $j = 0, 1, 2$.

Theorem 4.1. *An SLie (respectively W-) groupoid $\mathcal{G} \Rightarrow M$ with a chosen good chart (respectively good étale chart) G_0 of $\mathcal{G}$ corresponds to a Lie 2-groupoid (respectively 2-étale Lie 2-groupoid) $X_2 \Rightarrow X_1 \Rightarrow X_0$.*

4.1 The construction of $X_2 \Rightarrow X_1 \Rightarrow X_0$

Given an SLie groupoid $\mathcal{G} \Rightarrow M$, let $G_1 \Rightarrow G_0$ be a good groupoid presentation of $\mathcal{G}$ and E_m a bimodule presenting the morphism m . Let $J_l : E_m \rightarrow G_0 \times_M G_0$ and $J_r : E_m \rightarrow G_0$ be the moment maps of the bimodule E_m . Notice that for an SLie groupoid $g \cdot 1 = 1$ up to a 2-morphism, that is $m|_{\mathcal{G} \times_M M} = \text{id}$ up to a 2-morphism. Translating this into groupoid language, $J_1^{-1}(G_0 \times_M M)$ is the bimodule presenting $m|_{\mathcal{G} \times_M M}$. By the definition of SLie groupoids, there is an isomorphism $b_r : J_l^{-1}(G_0 \times_M M) \rightarrow G_1$. Similarly, there is an isomorphism $b_l : J_l^{-1}(M \times_M G_0) \rightarrow G_1$.

We construct

$$X_0 = M, X_1 = G_0, X_2 = E_m$$

with the structure maps

$$\begin{aligned} d_0^1 = \mathbf{s}, d_1^1 = \mathbf{t} : X_1 \rightarrow X_0, \quad d_0^2 = pr_2 \circ J_l, d_1^2 = J_r, d_2^2 = pr_1 \circ J_l : X_2 \rightarrow X_1, \\ s_0^0 = e : X_0 \rightarrow X_1, \quad s_0^1 = b_l^{-1} \circ e_G, s_1^1 = b_r^{-1} \circ e_G : X_1 \rightarrow X_2 \end{aligned} \quad (9)$$

where pr_i is the i -th projection $G_0 \times_M G_0 \rightarrow G_0$, $\mathbf{s}, \mathbf{t}$ present $\bar{\mathbf{s}}, \bar{\mathbf{t}} : \mathcal{G} \Rightarrow M$, and e_G is the identity embedding $G_0 \rightarrow G_1$. We still need the 3-multiplication maps


$$m_i : \Lambda(X)_{3,i} \rightarrow X_2 \quad i = 0, \dots, 3.$$

Let us first construct m_0 . Notice that in the 2-associative diagram, we have a 2-morphism $a : m \circ (m \times id) \rightarrow m \circ (id \times m)$. Translating this into the language of groupoid, we have the following isomorphism of bimodules:

$$a : ((E_m \times_{G_0} G_1) \times_{G_0 \times_M G_0} E_m) / (G_1 \times_M G_1) \rightarrow ((G_1 \times_{G_0} E_m) \times_{G_0 \times_M G_0} E_m) / (G_1 \times_M G_1). \quad (10)$$

Suppose $(\eta_1, \eta_2, \eta_3) \in \Lambda(X)_{3,0}$. Then $(\eta_3, 1, \eta_1)$ represents a class in $(E_m \times_{G_0} G_1) \times_{G_0 \times_M G_0} E_m / \sim$ (we write $\sim$ when it is clear which groupoid action is meant). Moreover, its image under a can be represented by $(1, \eta_0, \eta_2)$, that is,

$$a([(\eta_3, 1, \eta_1)]) = [(1, \eta_0, \eta_2)].$$



As before we imagine the j -dimensional faces of the picture are elements of X_j . Then we arrive naturally at η_0 . To show that the above definition is good, we have to show: 1) the image under a can be represented by some element of the form $(1, \eta_0, \eta_2)$; 2) the choice of η_0 is unique; 3) this map is smooth.

Suppose $a([(\eta_3, 1, \eta_1)]) = [(1, \tilde{\eta}_0, \tilde{\eta}_2)]$. Notice that the $G_1 \times_M G_1$ action on G_1 on both sides of (10) are the right multiplication by one of the copies of G_1 , therefore we can always suppose the element in G_1 is 1.

Notice that similarly to the proof of Lemma 8.1, the map $m \times pr_1 : \mathcal{G} \times_{\bar{\mathbf{s}}, M, \bar{\mathbf{t}}} \mathcal{G} \rightarrow \mathcal{G} \times_{\bar{\mathbf{t}}, M, \bar{\mathbf{t}}} \mathcal{G}$ is an isomorphism. Therefore $E_m \times_{pr_1 \circ J_l, G_0, \mathbf{t}_G} G_1$ is a Morita bibundle from the Lie groupoid $G_1 \times_{\mathbf{s}_G, M, \mathbf{t}_G} G_1 \rightrightarrows G_0 \times_{\mathbf{s}, M, \mathbf{t}} G_0$ to $G_1 \times_{\mathbf{t}_G, M, \mathbf{s}_G} G_1 \rightrightarrows G_0 \times_{\mathbf{t}, M, \mathbf{t}} G_0$, where $\mathbf{s}_G, \mathbf{t}_G$ are the source and target maps on G and $\mathbf{s}, \mathbf{t}$ are the maps $G_0 \rightarrow M$ presenting $\bar{\mathbf{s}}, \bar{\mathbf{t}}$. Here the two moment maps are J_l (for E_m) and $J_r \times \mathbf{s}_G$. Therefore the left groupoid action of $G_1 \times_{\mathbf{s}_G, M, \mathbf{t}_G} G_1$ is principal on the bibundle $E_m \times_{pr_1 \circ J_l, G_0, \mathbf{t}_G} G_1$. Notice that $(\eta_2, 1)$ and $(\tilde{\eta}_2, 1)$ are on the same fibre of $J_r \times \mathbf{s}_G$ over $G_0 \times_{\mathbf{t}, M, \mathbf{t}} G_0$ because the way we arrange the map makes them share the same edges $1 \rightarrow 0$ and $3 \rightarrow 0$. So there is a unique groupoid element (γ_1, γ_2) such that $(\tilde{\eta}_2, 1) \cdot (\gamma_1, \gamma_2) = (\eta_2, 1)$. This also forces γ_2 to be 1. Therefore, we have

$$(\tilde{\eta}_2, 1, \tilde{\eta}_0) \cdot (\gamma_1, 1) = (\eta_2, 1, \eta_0).$$

By uniqueness of γ_1 , the choice of η_0 is unique. Moreover, since smoothness is a local property and the map a is an isomorphism and the groupoid action is principal, our map is naturally smooth.

For other m 's, we can precede in a similar fashion. More precisely, for m_1 one can make the same definition for m_0 but using a^{-1} . It is even easier to define m_2 and m_3 . For

example, for m_2 , on the right hand side of (10), since the $G_1 \times_M G_1$ action on $G_1 \times_M E_m$ is principal, the analogous statement to 1) naturally holds. Thus we realize that given any three η 's, we can always put them in the same spots as we did for m_0 . Then any three of them determine the fourth. Hence the m 's are compatible with each other.

4.2 Proof that what we construct is a Lie 2-groupoid

By Prop-Def. 2.3, to show the above construction gives us a Lie 2-groupoid, we just have to show that the m 's satisfy the coherence conditions and associativity, and 1-Kan 2-Kan conditions. Condition 1-Kan is implied by the fact that $\mathbf{s}, \mathbf{t} : G_0 \Rightarrow M$ are surjective submersions; $Kan(2, 1)$ is implied by the fact that the moment map $J_l : E_m \rightarrow G_0 \times_{\mathbf{s}, M, \mathbf{t}} G_0$ is a surjective submersion; $Kan(2, 0)$ is implied by the surjective submersion

$$J_l : E_m \cong (G_1 \times_M E_i \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1 \rightarrow G_0 \times_{\mathbf{t}, M, \mathbf{t}} G_0 = \Lambda[2, 0](X),$$

where $E_i \cong G_1$ (Lemma 8.3) is the bimodule presenting the inverse map of $\mathcal{G}$, the composed bibundle is the one presenting the map $(g, h) \mapsto (g, h^{-1}) \mapsto gh^{-1}$; $Kan(2, 2)$ follows similarly as $Kan(2, 0)$.

The coherence conditions

The first identity in (5) corresponds to an identity of 2-morphisms,

$$(1 \cdot (g_1 \cdot g_2) \stackrel{a}{\sim} (1 \cdot g_1) \cdot g_2 \sim g_1 \cdot g_2) = (1 \cdot (g_1 \cdot g_2) \sim g_1 \cdot g_2),$$

Restrict the two bimodules in (10) to $M \times_M G_0 \times_M G_0$, then we get E_m on the left hand side because $J_l^{-1}(M \times_M G_0) = G_1$ and $((G_1 \times_M G_1) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1 = E_m$. More precisely, the elements in $(E_m \times_{G_0} G_1) \times_{G_0 \times_M G_0} E_m |_{M \times_M G_0 \times_M G_0} / \sim$ have the form $[(s_0 \circ d_2(\eta), 1, \eta)]$, and the isomorphism to E_m is given by $[(s_0 \circ d_2(\eta), 1, \eta)] \mapsto \eta$. Similarly for the right hand side, i.e. $[(s_0 \circ d_1(\eta), 1, \eta)] \mapsto \eta$ gives the other isomorphism. By 4c in Def. 3.1, the composition of the first and the inverse of the second map is a (restricted on the restricted bimodules), so we have

$$a([(s_0 \circ d_2(\eta), 1, \eta)]) = ([(1, \eta, s_0 \circ d_1(\eta))]),$$

which implies the first identity in (5). The rest follows similarly.

Associativity

For the associativity we will have to use the cube condition 3c for a in Def. 3.1 (called also “pentagon condition” in the literature). Let η_{ijk} denote the faces in X_2 fitting in diagram (6). Suppose we are given the faces $\eta'_{0i4}s \in X_2$ and the faces $\eta'_{0ij}s \in X_2$. Then we have two ways to determine the face η_{123} using m 's as described in Prop-Def. 2.3. We will show below that these two constructions give the same element in X_2 .

Translate the cube condition into the language of Lie groupoids. The morphisms become bimodules and the 2-morphisms become the morphisms between bimodules. The cube condition tells us that the following two compositions of morphisms are the same (here for

simplicity, we omit writing the base space of the fibre products and the groupoids by which we take quotients):

$$\begin{aligned}
(E_m \times G_1 \times G_1) \times (E_m \times G_1) \times E_m / \sim &<--> ((g_1 g_2) g_3) g_4 \\
\stackrel{id \times a}{\longrightarrow} (E_m \times G_1 \times G_1) \times (G_1 \times E_m) \times E_m / \sim &<--> (g_1 g_2) (g_3 g_4) \\
\stackrel{id}{\longrightarrow} (G_1 \times G_1 \times E_m) \times (E_m \times G_1) \times E_m / \sim &<--> (g_1 g_2) (g_3 g_4) \\
\stackrel{id \times a}{\longrightarrow} (G_1 \times G_1 \times E_m) \times (G_1 \times E_m) \times E_m / \sim &<--> g_1 (g_2 (g_3 g_4))
\end{aligned}$$

and

$$\begin{aligned}
(E_m \times G_1 \times G_1) \times (E_m \times G_1) \times E_m / \sim &<--> ((g_1 g_2) g_3) g_4 \\
\stackrel{a \times id}{\longrightarrow} (G_1 \times E_m \times G_1) \times (E_m \times G_1) \times E_m / \sim &<--> (g_1 (g_2 g_3)) g_4 \\
\stackrel{id \times a}{\longrightarrow} (G_1 \times E_m \times G_1) \times (G_1 \times E_m) \times E_m / \sim &<--> g_1 ((g_2 g_3) g_4) \\
\stackrel{a \times id}{\longrightarrow} (G_1 \times G_1 \times E_m) \times (G_1 \times E_m) \times E_m / \sim &<--> g_1 (g_2 (g_3 g_4))
\end{aligned}$$

Tracing through where the element $(\eta_{034}, (\eta_{023}, 1), (\eta_{012}, 1, 1))$ goes via the first and second composition, it should end up in the same element. So we have

$$\begin{aligned}
& [((\eta_{012}, 1, 1)), (\eta_{023}, 1), \eta_{034}] \\
& \stackrel{id \times a}{\mapsto} [((\eta_{012}, 1, 1), (1, \eta_{234}), \eta_{024})] \\
& \stackrel{id}{\mapsto} [((1, 1, \eta_{234}), (\eta_{012}, 1), \eta_{024})] \\
& \stackrel{id \times a}{\mapsto} [((1, 1, \eta_{234}), (1, \eta_{124}), \eta_{014})]
\end{aligned}$$

where $\eta_{234} = m_0(\eta_{034}, \eta_{024}, \eta_{023})$ and $\eta_{124} = m_0(\eta_{024}, \eta_{014}, \eta_{012})$;

$$\begin{aligned}
& [((\eta_{012}, 1, 1), (\eta_{023}, 1), \eta_{034})] \\
& \stackrel{a \times id}{\mapsto} [((1, \eta_{123}, 1), (\eta_{013}, 1), \eta_{034})] \\
& \stackrel{id \times a}{\mapsto} [((1, \eta_{123}, 1), (1, \eta_{134}), \eta_{014})] \\
& \stackrel{a \times id}{\mapsto} [((1, 1, \eta_{234}), (1, \eta_{124}), \eta_{014})]
\end{aligned}$$

where $\eta_{123} = m_0(\eta_{023}, \eta_{013}, \eta_{012})$ and $\eta_{134} = m_0(\eta_{034}, \eta_{014}, \eta_{013})$. Therefore, the last map tells us that

$$\eta_{123} = m_3(\eta_{234}, \eta_{134}, \eta_{124}).$$

Therefore associativity holds!

Comments on the étale condition

It is easy to see that if $G_1 \Rightarrow G_0$ is étale, by principality of the right G action on E_m , the map $E_m \rightarrow G_0 \times_M G_0$ is étale. Moreover since $E_m \rightarrow \Lambda(X)_{2,j} = \Lambda[2, j](X)$ is a surjective submersion by $Kan(2, j)$, by dimension counting, it is furthermore an étale map.

5 From Lie 2-groupoids to stacky Lie groupoids

If X is a Lie 2-groupoid, then $G_1 := d_2^{-1}(s_0(X_0)) \subset X_2$, which is the set of bigons, is a Lie groupoid over $G_0 := X_1$ (Lemma 5.2). Here we might notice that there is another natural choice for the space of bigons, namely $\tilde{G}_1 := d_0^{-1}(s_0(X_0))$. But $G_1 \cong \tilde{G}_1$ by the following observation: given an element $\eta_3 \in G_1$, it fits as the face opposite to 3 in diagram (11) with $1 \rightarrow 0$ and $2 \rightarrow 3$ degenerate and η_2, η_1 degenerate; then m_0 gives a morphism $\varphi : G_1 \rightarrow \tilde{G}_1$ and m_3 gives the inverse. Therefore we might consider only G_1 . Then $G_1 \Rightarrow G_0$ presents a stack which has an additional groupoid structure. Hence we have the statement from 2-groupoids to stacky Lie groupoids:


Theorem 5.1. *A Lie 2-groupoid (respectively 2-étale Lie 2-groupoid) X corresponds to an SLie (respectively W-) groupoid $\mathcal{G} \Rightarrow X_0$ where $\mathcal{G}$ is presented by the Lie groupoid $G_1 \Rightarrow G_0$.*

We prove this theorem by several lemmas.

5.1 The stack $\mathcal{G}$

Lemma 5.2. *$G_1 \Rightarrow G_0$ is a Lie groupoid.*

Proof. The target and source maps are given by d_0^1 and d_1^1 . The identity $G_0 \rightarrow G_1$ is given by $s_0^1 : X_1 \rightarrow X_2$. The image of s_0^1 is in $G_1 (\subset X_2)$. Their compatibility conditions are implied by the compatibility conditions of the structure maps of simplicial manifolds. This will in particular imply that the identity is an embedding and **s** and **t** are surjective submersions after we establish the multiplication. The multiplication is given by the 3-multiplication of X by the following picture:



where η_3 and its edges $1 \rightarrow 0$, $2 \rightarrow 0$ and $2 \rightarrow 1$ are degenerate (one should imagine them being very short). Let η_i be the face facing i . Then any $(\eta_0, \eta_2) \in G_1 \times_{s, G_0, t} G_1$ fits in the above picture. We define $\eta_0 \cdot \eta_2 = m_1(\eta_0, \eta_2, \eta_3)$ with η_3 is the degenerate face in $s_0 \circ s_0(X_0)$ corresponding to the point $0 (=1=2)$. Then the associativity of the 3-multiplications ensures the associativity of “ $\cdot$ ”. The inverse map is also given by 3-multiplications: $\eta_2^{-1} = m_0(\eta_1, \eta_2, \eta_3)$ with η_1 the degenerate face in $s_0^1(X_1)$. $\square$

Remark 5.3. Similar construction shows that $\tilde{G}_1 \Rightarrow G_0$ with **t** = d_2^1 , **s** = d_1^1 is a Lie groupoid isomorphic to $G_1 \Rightarrow G_0$ via the map φ^{-1} .

5.2 Proof that $\mathcal{G} \Rightarrow M$ is an SLie groupoid

Source target maps and multiplication

There are three maps $d_i^2 : X_2 \rightarrow X_1 = G_0$ and they (as the moment maps of the action) all correspond to a groupoid action respectively. The actions are similarly given by the

3-multiplications as the multiplication of G_1 . The axioms of the actions are given by the associativity. For example, for d_1^2 , any $(\eta_0, \eta_2) \in X_2 \times_{d_1^2, X_1, \mathbf{t}_G} G_1$ fits inside picture (11), but one imagines that only $1 \rightarrow 0$ is a short (η_3 degenerate) edge. Then

$$\eta_0 \cdot \eta_2 := m_1(\eta_0, \eta_2, s_0 d_2(\eta_0)). \quad (12)$$

Moreover, notice that the four ways to compose source target and face maps $G_1 \xrightarrow[\mathbf{t}_G]{\mathbf{s}_G} G_0 \xrightarrow[d_1^1]{d_0^1} X_0$ only give two different maps: $d_0^1 \mathbf{s}_G$ and $d_1^1 \mathbf{t}_G$. They are surjective submersions since d_i 's and $\mathbf{s}_G \mathbf{t}_G$ are so and they give the source and target maps $\bar{\mathbf{s}}, \bar{\mathbf{t}} : \mathcal{G} \Rightarrow X_0$ where $\mathcal{G}$ is the differentiable stack presented by $G_1 \Rightarrow G_0$. Therefore $\bar{\mathbf{s}}$ and $\bar{\mathbf{t}}$ are also surjective submersions (Lemma 4.2 in [TZ06a]). We use these two maps to form the product groupoid

$$G_1 \times_{d_0^1 \mathbf{s}_G, X_0, d_1^1 \mathbf{t}_G} G_1 \Rightarrow G_0 \times_{d_0^1, X_0, d_1^1} G_0 \quad (13)$$

which presents the stack $\mathcal{G} \times_{\bar{\mathbf{s}}, X_0, \bar{\mathbf{t}}} \mathcal{G}$. Then we have the following lemma:

Lemma 5.4. *$(X_2, d_2^2 \times d_1^2, d_1^2)$ is an H.S. bimodule from the groupoid in (13) to $G_1 \Rightarrow G_0$.*

Proof. By $\text{Kan}(2, 1)$, $d_2^2 \times d_1^2$ is a surjective submersion from X_2 to $G_0 \times_{d_0^1, X_0, d_1^1} G_0$, so we only have to show that the right action of $G_1 \Rightarrow G_0$ on X_2 is free and transitive. This is implied by $\text{Kan}(3, j)$ and $\text{Kan}(3, j)!$ respectively.


transitivity: any (η_1, η_0) such that $d_0^2(\eta_1) = d_0^2(\eta_0)$ and $d_2^2(\eta_0) = d_2^2(\eta_1)$ fits inside (11) with η_3 a degenerate face (only $1 \rightarrow 0$ the short edge). Then there exists $\eta_2 := m_2(\eta_0, \eta_1, \eta_3) \in G_1$, making $\eta_0 \cdot \eta_2 = \eta_1$.

freeness: if $(\eta_0, \eta_2) \in X_2 \times_{d_1, X_1, \mathbf{t}} G_1$ satisfies $\eta_0 \cdot \eta_2 (=m_1(\eta_0, \eta_2, \eta_3)) = \eta_0$. Then $\eta_2 = m_2(\eta_0, \eta_0, \eta_3)$ and η_3 is degenerate. Thus $m_2(\eta_0, \eta_0, \eta_3) = s_0^1(3 \rightarrow 1)$ is a degenerate face. Therefore $\eta_2 = 1$. $\square$

Therefore X_2 gives a morphism $m : \mathcal{G} \times_{X_0} \mathcal{G} \rightarrow \mathcal{G}$.

Lemma 5.5. *With the source and target maps constructed above, m is a multiplication of $\mathcal{G} \Rightarrow X_0$.*

Proof. By construction, it is clear that $\bar{\mathbf{t}} \circ m = \bar{\mathbf{t}} \circ pr_1$ and $\bar{\mathbf{s}} \circ m = \bar{\mathbf{s}} \circ pr_2$, where $pr_i : \mathcal{G} \times_{\bar{\mathbf{s}}, X_0, \bar{\mathbf{t}}} \mathcal{G} \rightarrow \mathcal{G}$ are the projections (see the left picture below).



To show the associativity, we reverse the argument in Section 4.1. There, we used the 2-morphism a to construct the 3-multiplications. Now we use the 3-multiplications and their associativity to construct a . Given the two H.S. bibundles presenting $m \circ (m \times id)$ and $m \circ (id \times m)$ respectively, we want to construct a map a as in (10), where $E_m = X_2$ and $M = X_0$. Given any element in $(X_2 \times_{G_0} G_1) \times_{G_0 \times_{X_0} G_0} X_2 / G_1 \times_{X_0} G_1$, as in Section 4.1, we can write it in the form of $[(\eta_3, 1, \eta_1)]$, with $(\eta_1, \eta_2, \eta_3) \in \text{hom}(\Lambda[3, 0], X)$ for some η_2 . Then we define

$$a([(\eta_3, 1, \eta_1)]) := [(1, m_0(\eta_1, \eta_2, \eta_3), \eta_2)].$$

As before, we need to show that: 1) the R.H.S. is indeed in $(G_1 \times_{G_0} X_2) \times_{G_0 \times_{X_0} G_0} X_2 / \sim$; 2) the definition of a does not depend on the choice of η_1 , η_3 and η_2 ; 3) a is smooth. The argument is the same. Here we only show 2) which is less obvious. First of all, if we choose a different $\tilde{\eta}_2$, since (η_1, η_2, η_3) and $(\eta_1, \tilde{\eta}_2, \eta_3)$ are both in $\text{hom}(\Lambda[3, 0], X)$, we have $d_2^2(\eta_2) = d_2^2(\tilde{\eta}_2)$ and $d_1^2(\eta_2) = d_1^2(\tilde{\eta}_2)$. So $\eta_2 = \eta_{013}$ and $\tilde{\eta}_2 = \eta_{01'3}$ form a degenerate horn (see the second picture of (14)). By $Kan(3, 0)$ there exists $\gamma = \gamma_{1'13} \in G_1$ such that $(1, \gamma) \cdot \tilde{\eta}_2 (= \gamma \cdot \tilde{\eta}_2) = \eta_2$, that is $\eta_{013} = m_1(\gamma, \eta_{01'3}, s_1^1(0 \rightarrow 1))$. Then by associativity and the definition of the G_1 action (12), we have $m_0(\eta_{023}, \eta_{01'3}, \eta_{01'2}) = \eta_{1'23} = \eta_{123} \cdot \gamma$. Therefore we have $[(1, \eta_{1'23}, \tilde{\eta}_2)] = [(1, \eta_{123}, \eta_2)]$. So the choice of η_2 will not affect the definition of a . If we choose a different $(\tilde{\eta}_3 = \eta_{0'12}, 1, \tilde{\eta}_1 = \eta_{0'23})$, then we can suppose $\eta_3 = \tilde{\eta}_3 \gamma_{00'2}$ and $\tilde{\eta}_1 = (\gamma_{00'2}, 1) \cdot \eta_1 = \gamma \cdot \eta_1$ (see the third picture of (14)). Then $(\tilde{\eta}_1, \eta_2, \tilde{\eta}_3) \in \text{hom}(\Lambda[3, 0], X)$ and

$$m_0(\tilde{\eta}_1, \eta_2, \tilde{\eta}_3) = \eta_{123} = m_0(\eta_1, \eta_2, \eta_3).$$

So this choice will not affect a neither.

Now the higher coherence of a follows from the associativity by the same argument as in Section 4.2. $\square$

Identity


Now we notice that $s_0 : X_0 \hookrightarrow G_0$ and $e_G \circ s_0 : X_0 \hookrightarrow G_1$ with e_G the identity of G form a groupoid morphism from $X_0 \rightrightarrows X_0$ to $G_1 \rightrightarrows G_0$. It gives a morphism $\bar{e} : X_0 \rightarrow \mathcal{G}$ on the level of stacks.

Lemma 5.6. *$\bar{e}$ is the identity of $\mathcal{G}$.*

Proof. Recall from Definition 3.1 that we need to show that there is a 2-morphism b_l between the two maps $m \circ (\bar{e} \times id)$ and $pr_2 : X_0 \times_{X_0, t} \mathcal{G} \rightarrow \mathcal{G}$, and similarly a 2-morphism b_r . In our case, the H.S. bibundles presenting these two maps are $X_2|_{X_0 \times_{X_0} G_0}$ and G_1 respectively and they are the same by construction, hence $b_l = id$. For b_r , we notice that $X_2|_{G_0 \times_{X_0} X_0} = \tilde{G}_1$ and the isomorphism $\varphi^{-1} : \tilde{G}_1 \rightarrow G_1$ is b_r .

Now we need to show the higher coherences that b 's satisfy. The proofs are similar and here we prove 4b in Definition 3.1 in detail. Translating the diagram (7) in the language of

groupoids and bibundles, we obtain



Corresponding to (8), we need to show that the following diagram commute:

$$\begin{array}{ccc}
 (X_2 \times_{X_0} X_0)_{G_0 \times_{X_0} X_0} \tilde{G}_1 / \sim & \xrightarrow{a} & (G_1 \times_{X_0} \tilde{G}_1)_{G_0 \times_{X_0} G_0} X_2 / \sim \xrightarrow{b_r} (G_1 \times_{X_0} G_1)_{G_0 \times_{X_0} G_0} X_2 / \sim \\
 \downarrow b_r & & \nearrow \cong \\
 (X_2 \times_{X_0} X_0)_{G_0 \times_{X_0} X_0} G_1 / \sim & \xrightarrow{\cong} & X_2
 \end{array}$$

$$\begin{array}{ccccc}
 [(\eta_3, 1, \eta_1)] & \longmapsto & [(1, \eta_0 = 1, \eta_2)] & \longmapsto & [(1, \eta'_0 = 1, \eta_2)] \\
 \downarrow & & & & \nearrow ? \\
 [(\eta_{012}, 1, \eta_{0'02})] & \longmapsto & \eta_{012} \cdot \eta_{0'02}^{-1} & &
 \end{array}$$

Let us explain the diagram: A $[(\eta_3, 1, \eta_1)] \in (X_2 \times_{X_0} X_0) \times_{G_0 \times_{X_0} X_0} \tilde{G}_1 / \sim$ fits inside (15) with $3 \rightarrow 2$ a degenerate edge and $\eta_0 = \eta_{123}$ a degenerate face⁹, therefore its image under a is $[(1, \eta_0 = 1, \eta_2)]$. By the construction of b_r , $\eta'_0 = b_r(\eta_0) = 1$ is also degenerate. Then $[(1, \eta'_0 = 1, \eta_2)]$ maps to η_2 under “ $\cong$ ”. Now we follow the other direction of the maps. Then $b_r(\eta_{023}) = \eta_{0'02}$. The image of $[(\eta_{012}, 1, \eta_{0'02})]$ under “ $\cong$ ” is $\eta_{012} \cdot \eta_{0'02}^{-1} = m_1(\eta_{0'02}^{-1}, \eta_{0'01} = 1, \eta_{012}) = \eta_{0'12}$. Notice that most of the “flat” faces are degenerate except for $\eta_{0'02}$ and η_{023} . Therefore consider the 3-simplices $(0', 0, 1, 3)$ and $(0', 1, 2, 3)$, then we have $\eta_{013} = \eta_{0'13}$ and $\eta_{0'13} = \eta_{0'12}$ by (5). Since η_{013} is η_2 , the desired diagram commutes. $\square$

⁹Now when it is confusing, we call a face by its three vertices, for example now $\eta_1 = \eta_{023}$.

Inverse

By Prop. 8.4, we only have to show that the actions of G and G^{op} on $X_2 \times_{d_1^2, G_0} M$, induced respectively by the first and second components of the left action of $G_1 \times_M G_1 \Rightarrow G_0 \times_M G_0$, are principal (see (21)). We prove this for the first copy of G_1 and the proof for the second is similar. A $(\eta_3, \eta_1) \in G_1 \times_{s_G, G_0, d_0^2} X_2 \times_{d_1^2, G_0} M$ fits inside (11) with η_2 a degenerate face corresponding to the point $0 = 1 = 3$. Then the freeness of the action is implied by $Kan!(3, 0)$ and the transitivity of the action is implied by $Kan(3, 0)$.

Comments on the étale condition

If $X_2 \rightarrow \Lambda[2, j](X)$ are étale maps, then the moment map $J_l : E_m \cong X_2 \rightarrow G_0 \times_{s, M, t} G_0$ is étale. By principality of the right action of $G_1 \Rightarrow G_0$, it is an étale groupoid. This concludes the proof of the Theorem 5.1.

6 Equivalences of Lie 2-groupoids

As in Theorem 4.1, from an SLie groupoid to a Lie 2-groupoid, one has to choose a chart of the stack $\mathcal{G}$. Therefore to construct a 1-1 correspondence between SLie groupoids and Lie 2-groupoids, we have to quotient by some sort of equivalence of 2-groupoids since one could have different 2-groupoids X and Y corresponding to different charts G_0 and H_0 of the same stack $\mathcal{G}$. In this section, we establish various equivalences of Lie 2-groupoids.

We first look at the more general case of Lie homotopy types and motivate our later definitions. The proof of this direction itself is carried out in a later work [HZ].

6.1 Strict maps of Lie homotopy types

The reader's first guess is probably that a morphism $f : X \rightarrow Y$ of Lie homotopy types ought to be a simplicial smooth map i.e. a collection of smooth maps $f_n : X_n \rightarrow Y_n$ that commute with faces and degeneracies. In the language of categories, this is just a natural transformation from the functor X to the functor Y . We shall call such a natural transformation a *strict map* from X to Y . Unfortunately, it is known that, already in the case of usual Lie groupoids, such strict notions are not good enough. Indeed there are strict maps that are not invertible even though they ought to be isomorphisms. That's why people introduced the notion of *Hilsum-Skandalis maps* [Mrč]. Here is an example of such a map: consider a manifold M with an open cover $\{\mathcal{U}_\alpha\}$. The simplicial manifold X with $X_n = \bigsqcup_{\alpha_1, \dots, \alpha_n} \mathcal{U}_{\alpha_1} \cap \dots \cap \mathcal{U}_{\alpha_n}$ maps naturally to the constant simplicial manifold M . All the fibers of that map are simplices, in particular they are contractible simplicial sets. Nevertheless, that map has no inverse.

The second guess is then to define a special class of strict maps which we shall call *equivalences*. A map from X to Y would then be a *zig-zag* of strict maps $X \xleftarrow{\sim} Z \rightarrow Y$, where the map $Z \rightarrow X$ is one of these equivalences.

This will not be our final choice for what a morphism of Lie homotopy types should be. The notion of equivalence is nevertheless very useful (e.g. to define sheaf cohomology of Lie homotopy types) and we will study it in this section. It is very much inspired from the notion of equivalence of simplicial sets (sometimes called weak equivalence). Recall ([GJ99])

that a map $S \rightarrow T$ of simplicial sets is an equivalence if it induces isomorphisms of all the topological homotopy groups. Here is an equivalent condition which we will generalize to Lie homotopy types:

Lemma 6.1. *A map $S \rightarrow T$ of Kan simplicial sets is an equivalence if, for any $m \geq 0$ and any commutative solid arrow diagram*

$$\begin{array}{ccc} \partial\Delta^n & \longrightarrow & S \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^n & \longrightarrow & T \end{array} \quad (16)$$

there exists a dotted arrow that makes both triangles commute. Here $\partial\Delta^n$ stands for the boundary of the n -simplex.

Proof. Let $\alpha \in \pi_{n-1}(S)$ be represented by some map $\partial\Delta^n \rightarrow S$ and assume $\alpha \mapsto 0 \in \pi_{n-1}(T)$. This means that we have a map $\Delta^n \rightarrow T$ that makes diagram (16) commute. By hypothesis, we get a map $\Delta^n \rightarrow S$ therefore $\alpha = 0 \in \pi_{n-1}(S)$. This proves the injectivity of $\pi_{n-1}(S) \rightarrow \pi_{n-1}(T)$.

Now let us consider an element $\beta \in \pi_n(T)$ represented by a map $(\Delta^n, \partial\Delta^n) \rightarrow (T, *)$. That map fits into a diagram (16) where the top arrow sends everything to the base point. By hypothesis we get a map $(\Delta^n, \partial\Delta^n) \rightarrow (S, *)$. It represents an element of $\pi_n(S)$ that is mapped to β . This proves the surjectivity of $\pi_n(S) \rightarrow \pi_n(T)$.

The Kan condition has been used implicitly to find representatives of the various homotopy classes (otherwise we would have needed to subdivide the simplices). $\square$

Translating the condition of lemma 6.1 into hom spaces gives:

Definition 6.2. A strict map $f : Z \rightarrow X$ of Lie homotopy types is an equivalence if the natural maps from $Z_n = \text{hom}(\Delta^n, Z)$ to the pull-back $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow \text{hom}(\Delta[n], X))$ are surjective submersions for all $n \geq 0$.

The proof of this being an equivalence is in [HZ]. But even before this, we will need to talk a lot about spaces of the form $PB(\text{hom}(A, Z) \rightarrow \text{hom}(A, X) \leftarrow \text{hom}(B, X))$, where the maps are induced by some fixed maps $A \rightarrow B$ and $Z \rightarrow X$. To avoid the cumbersome pull-back notation, we shall denote these spaces by

$$\left\{ \begin{array}{ccc} A & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ B & \xrightarrow{?} & X \end{array} \right\}.$$

This notation indicates that the space parameterizes all commuting diagrams of the form

$$\begin{array}{ccc} A & \longrightarrow & Z \\ \downarrow & & \downarrow \\ B & \longrightarrow & X, \end{array}$$

where we allow the horizontal arrows to vary but we fix the vertical ones.

As in the case of definition 1.2, we need to justify why the pull-backs used in definition 6.2 are manifolds. This is specially surprising since the spaces $\text{hom}(\partial\Delta^n, Z)$ need not be manifolds (for example take $n = 2$ and Z the cross product groupoid associated to the action of S^1 on $\mathbb{R}^2$ by rotation around the origin).

Lemma 6.3. *Let S be a finite collapsible simplicial set and $T \rightarrow S$ a sub-simplicial set of dimension $\leq k$. Let $f : Z \rightarrow X$ be a strict map of Lie homotopy types such that the natural map*

$$Z_n \rightarrow \left\{ \begin{array}{ccc} \partial\Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{?} & X \end{array} \right\}$$

is a surjective submersion for all $n \leq k$ (assuming the right hand sides are known to be manifolds). Then the space

$$\left\{ \begin{array}{ccc} T & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\}$$

is naturally a manifold.

Proof. Let T' be a sub-simplicial set obtained by deleting one simplex from T (without its boundary). We have a push-out diagram

$$\begin{array}{ccc} T' & \xrightarrow{\quad} & T \\ \uparrow & \lrcorner & \uparrow \\ \partial\Delta^n & \hookrightarrow & \Delta^n \end{array} .$$

Similarly to the proof of lemma 2.1 this gives a pull-back diagram

$$\begin{array}{ccc} \left\{ \begin{array}{ccc} T' & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} T & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} \\ \downarrow & \lrcorner & \downarrow \\ \left\{ \begin{array}{ccc} \partial\Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} \Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} , \end{array}$$

which may be combined with the pull-back diagram

$$\begin{array}{ccc}
\left\{ \begin{array}{ccc} \partial\Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} \Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} \\
\downarrow & \lrcorner & \downarrow \\
\left\{ \begin{array}{ccc} \partial\Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} \Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{?} & X \end{array} \right\} = Z_n
\end{array}$$

to give yet another pull-back diagram

$$\begin{array}{ccc}
\left\{ \begin{array}{ccc} T' & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} T & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ S & \xrightarrow{?} & X \end{array} \right\} \\
\downarrow & \lrcorner & \downarrow \\
\left\{ \begin{array}{ccc} \partial\Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{?} & X \end{array} \right\} & \longleftarrow & \left\{ \begin{array}{ccc} \Delta^n & \xrightarrow{?} & Z \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{?} & X \end{array} \right\} = Z_n.
\end{array} \tag{17}$$

By induction on the size of T , and by lemma 2.1 (case $T = \emptyset$), we may assume that the upper left and lower left spaces in (17) are known to be manifolds. The bottom arrow is a submersion by hypothesis. Therefore by transversality, the upper right space is also a manifold, which is what we wanted to prove. $\square$

6.2 Morita Equivalence of Lie 2-groupoids

Now we define equivalences of Lie 2-groupoids based on Section 6.1. To simplify our notation, $\rightarrow$ and $\leftarrow$ always denote surjective submersions.

Definition 6.4. Given two Lie 2-groupoids $Z_2 \rightrightarrows Z_1 \rightrightarrows Z_0$ and $X_2 \rightrightarrows X_1 \rightrightarrows X_0$, a strict morphism f between them is an equivalence if the natural map from Z_n to the pull-back $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ are surjective submersions for all $n = 0, 1$ and an isomorphism for $n = 2$.

As in the case of Lie homotopy types, we find that $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ is a manifold.

Similarly to the pull-back Lie groupoid, we make the following observation: Let X be a 2-groupoid and $Z_1 \rightrightarrows Z_0$ be two manifolds with structure maps as in (1) up to the level $n \leq 1$, and $f_n : Z_n \rightarrow X_n$ preserving the structure maps d_k^n 's and s_k^{n-1} 's for $n \leq 1$. Then $\text{hom}(\partial\Delta^n, Z)$ still makes sense for $n \leq 1$. We further suppose $f_0 : Z_0 \rightarrow X_0$ (hence

$Z_0 \times Z_0 \times_{X_0 \times X_0} X_1$ is a manifold) and $Z_1 \rightrightarrows Z_0 \times Z_0 \times_{X_0 \times X_0} X_1$. That is to say that the induced map from Z_n to the pull-back $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ are surjective submersions for $n = 0, 1$. Then we form

$$Z_2 = PB(\text{hom}(\partial\Delta^2, Z) \rightarrow \text{hom}(\partial\Delta^2, X) \leftarrow X_2).$$


It is easy to see that the proof of Lemma 6.3 still guarantees that Z_2 a manifold. Moreover there are $d_i^2 : Z_2 \rightarrow Z_1$ induced by the natural projections $\text{hom}(\partial\Delta^2, Z) \rightarrow Z_1$; $s_i^1 : Z_1 \rightarrow Z_2$ by

$$s_0^1(h) = (h, h, s_0^0(d_0^1(h)), s_0^1(f_1(h))), \quad s_1^1(h) = (s_0^0(d_1^1(h)), h, h, s_1^1(f_1(h)));$$

$m_i : \Lambda(Z)_{3,i} \rightarrow Z_2$ by for example

$$m_0((h_2, h_5, h_3, \bar{\eta}_1), (h_4, h_5, h_0, \bar{\eta}_2), (h_1, h_3, h_0, \bar{\eta}_3)) = (h_2, h_4, h_1, m_0(\bar{\eta}_1, \bar{\eta}_2, \bar{\eta}_3)),$$

and similarly for other m 's.



Then $Z_2 \rightrightarrows Z_1 \rightrightarrows Z_0$ is a Lie 2-groupoid and we call it the *pull-back 2-groupoid* by f . Moreover $f : Z \rightarrow X$ is an equivalence with the natural projection $f_2 : Z_2 \rightarrow X_2$.

Definition 6.4 can be easily generalized to higher dimensions.

Definition 6.5. A strict map $f : Z \rightarrow X$ of Lie m -groupoids is an equivalence if the natural maps from Z_n to the pull-back $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ are surjective submersions for all $0 \leq n \leq m - 1$ and isomorphism for $n = m$.

Remark 6.6. First of all, it is easy to see that when $m = 1$, we get the equivalence (or pull-back) of Lie groupoids. Moreover, since for a Lie m -groupoid layers higher or equal to $m + 1$ are determined by layers lower or equal to m , the maps from Z_n to the pull-back $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ are isomorphisms for all $n \geq m$.

Similar to the proof that $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n)$ is a manifold, we have the following lemma:

Lemma 6.7. *The projection $PB(\text{hom}(\partial\Delta^n, Z) \rightarrow \text{hom}(\partial\Delta^n, X) \leftarrow X_n) \rightarrow X_n$ is a surjective submersion.*

Proof. We can use the same induction as in Lemma 6.3 and only have to notice that the upper lever map in (17) is a surjective submersion (then let $T' = \emptyset$). $\square$

Lemma 6.8. *The composition of equivalences is still an equivalence.*

Proof. Suppose that $Z \xrightarrow{f} Y \xrightarrow{g} X$ are two equivalences. Then we have

$$\begin{array}{ccccc}
Z_n \rightrightarrows \mathrm{hom}(\partial\Delta^n, Z) \times_{\mathrm{hom}(\partial\Delta^n, Y)} Y_n & \longrightarrow & Y_n & & \\
\downarrow \leftarrow & & \downarrow & & \\
\mathrm{hom}(\partial\Delta^n, Z) \times_{\mathrm{hom}(\partial\Delta^n, X)} X_n \rightrightarrows \mathrm{hom}(\partial\Delta^n, Y) \times_{\mathrm{hom}(\partial\Delta^n, X)} X_n & \longrightarrow & X_n & & \\
\downarrow & & \downarrow & & \downarrow \\
\mathrm{hom}(\partial\Delta^n, Z) & \longrightarrow & \mathrm{hom}(\partial\Delta^n, Y) & \longrightarrow & \mathrm{hom}(\partial\Delta^n, X)
\end{array}$$

Therefore we have $Z_n \rightrightarrows \mathrm{hom}(\partial\Delta^n, Z) \times_{\mathrm{hom}(\partial\Delta^n, X)} X_n$ when $n \leq 1$. When $n = 2$ one just has to replace $\rightrightarrows$ by $\cong$ in the above diagram and the result. $\square$

Remark 6.9. It is not hard to see this result generalizes to n -groupoids.

Lemma 6.10. *If $f : Z \rightarrow X$ and $f' : Z \rightarrow X$ are equivalences, then there exist a Lie 2-groupoid $Z'' = Z \times_X Z'$ and two equivalences $Z \leftarrow Z'' \rightarrow Z'$.*

Proof. First of all we prove that $Z''_n = Z_n \times_{X_n} Z'_n$ (a manifold by Lemma 6.7) with product face and degeneracy maps is a Lie 2-groupoid. We use Prop-Def. 2.3. Notice that the maps on both sides of $Z_0 \leftarrow Z_0 \times_{X_0} X_1 = PB(Z_0 \rightrightarrows X_0 \leftarrow X_1) \rightarrow X_1$ are surjective submersions by the property of pull-backs. Similarly we have a surjective submersion,

$$Z_0 \times Z_0 \times_{X_0 \times X_0} X_1 = PB((Z_0 \times_{X_0} X_1) \rightrightarrows X_1 \leftarrow (Z_0 \times_{X_0} X_1)) \rightarrow (Z_0 \times_{X_0} X_1),$$

hence $Z_1 \rightarrow Z_0 \times_{X_0} X_1$ by composing with $Z_1 \rightarrow Z_0 \times Z_0 \times_{X_0 \times X_0} X_1$. Similarly we have this for Z' . For the 1-Kan conditions of Z'' , we should show that the degeneracy maps $Z_1 \times_{X_1} Z'_1 \rightarrow Z''_0 = Z_0 \times_{X_0} Z'_0$ are surjective submersions. The degeneracy map $d_0^1 \times d_0^1$ is a surjective submersion by chasing the arrows denoted with “!” in the following diagram,

$$\begin{array}{ccccccc}
Z_1 \times_{X_1} Z'_1 & \xrightarrow{\quad} & * & \xrightarrow{\quad} & Z'_1 & & \\
\downarrow ! & \swarrow \leftarrow & \downarrow \leftarrow & \searrow \leftarrow & \downarrow & & \\
* & \xrightarrow{\quad} & * & \xrightarrow{\quad} & Z'_0 \times_{X_0} X_1 & & \\
\downarrow ! & \uparrow \leftarrow & \downarrow \leftarrow & \searrow \leftarrow & \downarrow & & \\
Z_1 & \xrightarrow{\quad} & Z_0 \times_{X_0} X_1 & \xrightarrow{\quad} & X_1 & & \\
& & \downarrow & & \downarrow & & \\
& & Z_0 \times_{X_0} Z'_0 & \xrightarrow{\quad} & Z'_0 & & \\
& & \downarrow & & \downarrow & & \\
& & Z_0 & \xrightarrow{\quad} & X_0 & & \\
& & & & \downarrow d_0 & &
\end{array}$$

where $*$ denotes non-important pull-backs. The same diagram proves the result for $d_1^1 \times d_1^1$ also. This argument in fact shows the following general result about surjective submersions of manifolds,

Claim 6.11. *If $U \rightrightarrows \bar{U}$, and the map $V \rightarrow \bar{V}$ factors through a surjective submersion $V \rightrightarrows \bar{V} \times_{\bar{U}} U$, and $W \rightarrow \bar{W}$ factors through $W \rightrightarrows U \times_{\bar{U}} \bar{W}$, then the natural map $V \times_U W \rightrightarrows \bar{V} \times_{\bar{U}} \bar{W}$ is a surjective submersion. Here we suppose all the fibre products to be manifolds.*

For the 2-Kan condition of Z'' , we need to show that the maps $Z_2'' \rightarrow \text{hom}(\Lambda[2, j], Z'')$ are surjective submersions. Since $\text{hom}(\Lambda[2, j], Z'') = \text{hom}(\Lambda[2, j], Z) \times_{\text{hom}(\Lambda[2, j], X)} \text{hom}(\Lambda[2, j], Z')$, by Claim 6.11 and $X_2 \twoheadrightarrow \text{hom}(\Lambda[2, j], X)$, we only have to show $Z_2 \twoheadrightarrow \text{hom}(\Lambda[2, j], Z) \times_{\text{hom}(\Lambda[2, j], X)} X_2$ and the similar statement for Z' . Notice that

$$Z_0 \times Z_0 \twoheadrightarrow X_0 \times X_0, \quad Z_1 \twoheadrightarrow Z_0 \times Z_0 \times_{X_0 \times X_0} X_1,$$

by assumption. Therefore

$$\begin{aligned} \text{hom}(\partial\Delta^2, Z) &= \text{hom}(\Lambda[2, j], Z) \times_{Z_0 \times Z_0} Z_1 \\ &\twoheadrightarrow \text{hom}(\Lambda[2, j], Z) \times_{Z_0 \times Z_0} Z_0 \times Z_0 \times_{X_0 \times X_0} X_1 \\ &= \text{hom}(\Lambda[2, j], Z) \times_{X_0 \times X_0} X_1 \\ &= \text{hom}(\Lambda[2, j], Z) \times_{\text{hom}(\Lambda[2, j], X)} \text{hom}(\partial\Delta^2, X), \end{aligned}$$

where in the last step we use again $\text{hom}(\partial\Delta^2, X) = \text{hom}(\Lambda[2, j], X) \times_{X_0 \times X_0} X_1$. Hence

$$\begin{aligned} Z_2 &= \text{hom}(\partial\Delta^2, Z) \times_{\text{hom}(\partial\Delta^2, X)} X_2 \\ &\twoheadrightarrow \text{hom}(\Lambda[2, j], Z) \times_{\text{hom}(\Lambda[2, j], X)} \text{hom}(\partial\Delta^2, X) \times_{\text{hom}(\partial\Delta^2, X)} X_2 \\ &= \text{hom}(\Lambda[2, j], Z) \times_{\text{hom}(\Lambda[2, j], X)} X_2. \end{aligned}$$

The 3-multiplications on Z_2'' are induced by those of X_2 , by the second row of the equation,

$$\begin{aligned} Z_2'' &= \text{hom}(\partial\Delta^2, Z) \times_{\text{hom}(\partial\Delta^2, X)} X_2 \times_{X_2} \text{hom}(\partial\Delta^2, Z') \times_{\text{hom}(\partial\Delta^2, X)} X_2 \\ &= \text{hom}(\partial\Delta^2, Z) \times_{\text{hom}(\partial\Delta^2, X)} X_2 \times_{\text{hom}(\partial\Delta^2, X)} \text{hom}(\partial\Delta^2, Z') \\ &= Z_2 \times_{\text{hom}(\partial\Delta^2, X)} \text{hom}(\partial\Delta^2, Z') \\ &= \text{hom}(\partial\Delta^2, Z'') \times_{\text{hom}(\partial\Delta^2, Z)} Z_2. \end{aligned} \tag{18}$$

In the last row, we use $\text{hom}(\partial\Delta^2, Z'') = \text{hom}(\partial\Delta^2, Z) \times_{\text{hom}(\partial\Delta^2, X)} \text{hom}(\partial\Delta^2, Z')$. Therefore, the 3-multiplications of Z_2'' satisfy all the axioms as they are essentially the 3-multiplications of X_2 . Now we have proven that Z'' is a Lie 2-groupoid. Moreover, from the last row, we can also see that the projection $Z_2'' \rightarrow Z_2$ preserves the 3-multiplications since both of theirs are induced by those of X_2 . An analogous result holds for Z' . Therefore to complete the proof, we only have to show

$$\begin{aligned} Z_1'' &= Z_1 \times_{X_1} Z_1' \twoheadrightarrow \text{hom}(\partial\Delta^1, Z'') \times_{\text{hom}(\partial\Delta^1, Z)} Z_1 \\ &= \text{hom}(\partial\Delta^1, Z') \times_{\text{hom}(\partial\Delta^1, X)} Z_1, \end{aligned}$$

since $\text{hom}(\partial\Delta^1, Z'') = \text{hom}(\partial\Delta^1, Z) \times_{\text{hom}(\partial\Delta^1, X)} \text{hom}(\partial\Delta^1, Z')$. But this can be easily deduced from the following pull-back diagram,

$$\begin{array}{ccc} Z_1 \times_{X_1} Z_1' & \xrightarrow{\quad} & Z_1' \\ \downarrow \leftarrow & & \downarrow \\ Z_1 \times_{\text{hom}(\partial\Delta^1, X)} \text{hom}(\partial\Delta^1, Z') & \longrightarrow & \text{hom}(\partial\Delta^1, Z') \times_{\text{hom}(\partial\Delta^1, X)} X_1 \\ \downarrow & & \downarrow \\ Z_1 & \xrightarrow{\quad} & X_1, \end{array}$$

and the fact that $Z_1' \twoheadrightarrow \text{hom}(\partial\Delta^1, Z') \times_{\text{hom}(\partial\Delta^1, X)} X_1$. □

Definition 6.12. Two Lie 2-groupoids X and Y are *Morita equivalent* if there is a Lie 2-groupoid Z such that both of the maps $X \leftarrow Z \rightarrow Y$ are equivalences. With the above two lemmas, this definition does give an equivalence relation. We call it *Morita equivalence* of Lie 2-groupoids.

6.3 1-Morita Equivalence of Lie 2-groupoids

However Morita equivalent Lie 2-groupoids correspond to Morita equivalent SLie groupoids [BZ]. Hence to obtain isomorphic SLie groupoids, we need a stricter notion of equivalence of Lie 2-groupoids.

Proposition-Definition 6.13. A strict map of 2-groupoids $f : Z \rightarrow X$ is a *1-equivalence* if it is an equivalence with f_0 an isomorphism. Two 2-groupoids X and Y are 1-Morita equivalent if there is a 2-groupoid Z such that both of the maps $X \leftarrow Z \rightarrow Y$ are 1-equivalences. It gives an equivalence relation between Lie 2-groupoids and we call it **1-Morita equivalence**.

Proof. It is easy to see that the composition of 1-equivalences is still a 1-equivalence. We just have to notice that in Lemma 6.10 if both equivalences are 1-equivalences, then the projections $Z_0 \leftarrow Z_0 \times_{X_0} Z'_0 \rightarrow Z'_0$ are isomorphisms since $Z_0 \times_{X_0} Z'_0 \cong Z_0 \cong X_0 \cong Z'_0$. $\square$

Remark 6.14. For a 1-equivalence $Z \rightarrow X$, since $f_0 : Z_0 \cong X_0$, we have $\text{hom}(\partial\Delta^1, Z) = \text{hom}(\partial\Delta^1, X)$. So the condition on f_1 in Definition 6.4 becomes $f_1 : Z_1 \rightarrow X_1$.

7 One-to-one correspondence between Lie 2-groupoids and SLie groupoids

In this section, we use two lemmas to prove the following theorem:

Theorem 7.1. *There is a 1-1 correspondence of Lie 2-groupoids (respectively 2-étale Lie 2-groupoids) modulo 1-Morita equivalence and SLie groupoids (respectively W-groupoids).*

W-groupoids are isomorphic if and only if they are isomorphic as SLie groupoids, and 1-Morita equivalent 2-étale Lie 2-groupoids are 1-Morita equivalent Lie 2-groupoids. Therefore the étale version of the theorem is implied by the general case and we only have to prove the general case.

For the lemma below, we fix our notation: X and Y are Lie 2-groupoids in the sense of Prop-Def. 2.3; $G_0 = X_1$ and $H_0 = Y_1$; $X_0 = Y_0 = M$. G_1 and H_1 are the spaces of bigons in X and Y respectively, that is $d_2^{-1}(s_0(Y_0))$; Both $G_1 \Rightarrow G_0$ and $H_1 \Rightarrow H_0$ are Lie groupoids and they present differentiable stacks $\mathcal{G}$ and $\mathcal{H}$ respectively. Moreover $\mathcal{G} \Rightarrow M$ and $\mathcal{H} \Rightarrow M$ are stacky groupoids.

Lemma 7.2. *If $f : Y \rightarrow X$ is an equivalence, then*

1. *the groupoid $H_1 \Rightarrow H_0$ constructed from Y satisfies $H_1 \cong G_1 \times_{G_0 \times_M G_0} H_0 \times H_0$ with the pull-back groupoid structure (therefore $\mathcal{G} \cong \mathcal{H}$);*
2. *the above map $\mathcal{G} \cong \mathcal{H}$ induces an SLie groupoid isomorphism, that is*

- (a) there are a 2-morphism $a : m_{\mathcal{G}} \rightarrow m_{\mathcal{H}} : \mathcal{G} \times_M \mathcal{G} (\cong \mathcal{H} \times_M \mathcal{H}) \rightarrow \mathcal{G} (\cong \mathcal{H})$ and a 2-morphism $b : \bar{e}_{\mathcal{G}} \rightarrow \bar{e}_{\mathcal{H}} : M \rightarrow \mathcal{G} \cong \mathcal{H}$;
- (b) between maps $\mathcal{G} \times_M \mathcal{G} \times_M \mathcal{G} \rightarrow \mathcal{G}$, there is a commutative diagram of 2-morphisms,

$$\begin{array}{ccc} m_{\mathcal{G}} \circ (m_{\mathcal{G}} \times id) & \xrightarrow{a_G} & m_{\mathcal{G}} \circ (id \times m_{\mathcal{G}}) \\ \downarrow a & & \downarrow a \\ m_{\mathcal{H}} \circ (m_{\mathcal{H}} \times id) & \xrightarrow{a_H} & m_{\mathcal{H}} \circ (id \times m_{\mathcal{H}}) \end{array}$$

where by abuse of notation a denotes the 2-morphisms generated by a such as $a \odot (a \times id)$;

- (c) between maps $M \times_M \mathcal{G} \rightarrow \mathcal{G}$ and maps $\mathcal{G} \times_M M \rightarrow \mathcal{G}$ there are commutative diagrams of 2-morphisms,

$$\begin{array}{ccc} m_{\mathcal{G}} \circ (\bar{e}_{\mathcal{G}} \times id) & \xrightarrow{b_l^G} & pr_2 \\ \downarrow a \odot b & \nearrow b_l^H & \\ m_{\mathcal{H}} \circ (\bar{e}_{\mathcal{H}} \times id) & & \end{array} \quad \begin{array}{ccc} m_{\mathcal{G}} \circ (id \times \bar{e}_{\mathcal{G}}) & \xrightarrow{b_r^G} & pr_1 \\ \downarrow a \odot b & \nearrow b_r^H & \\ m_{\mathcal{H}} \circ (id \times \bar{e}_{\mathcal{H}}) & & \end{array}$$

Proof. Since $Y_2 \cong \text{hom}(\partial\Delta^2, Y) \times_{\text{hom}(\partial\Delta^2, X)} X_2$, we have

$$\begin{aligned} H_1 &= d_2^{-1}(s_0(Y_0)) = d_2^{-1}(s_0(X_0)) \times_{d_1 \times d_0, X_1 \times_M X_1} Y_1 \times_M Y_1 \\ &= G_1 \times_{\mathbf{t}_G \times \mathbf{s}_G, G_0 \times_M G_0} H_0 \times_M H_0 \\ &= G_1 \times_{\mathbf{t}_G \times \mathbf{s}_G, G_0 \times G_0} H_0 \times H_0, \end{aligned}$$

where $\mathbf{t}_G, \mathbf{s}_G$ denote the target and source map of $G_1 \Rightarrow G_0$, and the last step follows from the fact that $(\mathbf{t}_G \times \mathbf{s}_G)(G_1) \subset G_0 \times_M G_0$ and f preserves simplicial structures. The multiplication on H_1 (respectively G_1) is given by 3-multiplications on Y_2 (respectively X_2). Therefore H has the pull-back groupoid structure since Y is the pull-back of X . So item (1) is proven.

To prove 2a, we translate it to the following groupoid diagram:

$$\begin{array}{ccccc} & H_1 \times_M H_1 & & E_{m_{\mathcal{H}}} = Y_2 & \\ & \Downarrow & & \swarrow & \searrow \\ G_1 \times_M G_1 & H_0 \times_M H_0 & \xleftarrow{E_{m_{\mathcal{G}}} = X_2} & & G_1 \times_M H_0 \\ \Downarrow & \swarrow & \searrow & & \Downarrow \\ G_0 \times_M G_0 & & & & G_0 \end{array}$$

We need to show that the following compositions of bibundles are isomorphic

$$\begin{aligned} (Y_2 \times_{G_0} G_1) / H_1 &= (Y_2 \times_{H_0} H_0 \times_{G_0} G_1) / H_1 \\ &\stackrel{a}{\cong} H_0 \times_M H_0 \times_{G_0 \times_M G_0} X_2 = (H_0 \times_M H_0 \times_{G_0 \times_M G_0} G_1 \times_M G_1 \times_{G_0 \times_M G_0} X_2) / G_1 \times_M G_1. \end{aligned} \tag{19}$$

By item (1), any element in $(Y_2 \times_{G_0} G_1)/H_1$ can be written as $[(\eta, 1)]$ with $\eta \in Y_2$, and we construct a by $[(\eta, 1)] \mapsto (d_2(\eta), d_0(\eta), f_2(\eta))$. First of all, we need to show that a is well-defined. For this we only have to notice the identification $\eta = (\bar{\eta}, h_0, h_1, h_2)$ with $\bar{\eta} = f_2(\eta) \in X_2$ and $h_i = d_i(\eta)$ since f_2 preserves degeneracy maps. Also $\gamma \in H_1$ can be written as $\gamma = (\bar{\gamma}, h_1, h'_1)$ with $\bar{\gamma} = f_2(\gamma) \in G_1$, then the H_1 action on Y_2 is induced in the following way,

$$(\bar{\eta}, h_0, h_1, h_2) \cdot (\bar{\gamma}, h_1, h'_1) = (\bar{\eta} \cdot \bar{\gamma}, h_0, h'_1, h_2)$$


where $h_i = d_i(\eta)$. Hence if $(\eta', 1) = (\eta, 1) \cdot (\bar{\gamma}, h_1, h'_1)$, then $\bar{\gamma} = 1$ and $a([(\eta', 1)]) = (h_2, h_0, \bar{\eta}) = a([(\eta, 1)])$. Given $(h_2, h_0, \bar{\eta}) \in H_0 \times_M H_0 \times_{G_0 \times_M G_0} X_2$, take any h_1 such that $f_1(h_1) = d_1(\bar{\eta})$, then $(h_0, h_1, h_2) \in \text{hom}(\partial \Delta^2, Y)$. Thus we construct a^{-1} by $(h_2, h_0, \bar{\eta}) \mapsto [((\bar{\eta}, h_0, h_1, h_2), 1)]$. By the action of H_1 it is easy to see that a^{-1} is also well-defined. For the 2-morphism b the proof is much easier since in this case all the H.S. morphisms are strict morphisms of groupoids. So we only have to notice the commutative diagram

$$\begin{array}{ccc} M & \longrightarrow & G_0 \\ \downarrow & \nearrow & \\ H_0 & & \end{array}$$


Recall that the 3-multiplications on Y_2 are induced from those of X_2 in the following form:

$$m_0((\bar{\eta}_1, h_2, h_5, h_3), (\bar{\eta}_2, h_4, h_5, h_0), (\bar{\eta}_3, h_1, h_3, h_0)) = (m_0(\bar{\eta}_1, \bar{\eta}_2, \bar{\eta}_3), h_2, h_4, h_1),$$

and similarly for other m 's.



We also translate 2b to diagrams of groupoids, where we denote n -product $\square_i \times_M \cdots \times_M \square_i$ by $\square_i^n$,



Then we have a diagram of 2-morphisms between composed bibundles

$$\begin{array}{ccc}
((Y_2 \times_M H_1 \times_{H_0^2} Y_2/H_1^2) \times_{G_0} G_1)/H_1 & \xrightarrow{a_H} & ((H_1 \times_M Y_2 \times_{G_0^2} Y_2)/H_1^2) \times_{G_0} G_1/H_1 \\
\downarrow a & & \downarrow a \\
(H_0^3 \times_{G_0^3} (X_2 \times_M G_1 \times_{G_0^2} X_2))/G_1^2 & \xrightarrow{a_G} & ((H_0^3 \times_{G_0^3} (G_1 \times_M X_2 \times_{G_0^2} X_2))/G_1^2)
\end{array}$$

which is

$$\begin{array}{ccc}
[(\eta_3, 1, \eta_2, 1)] & \xrightarrow{a_H} & [(1, \eta_0, \eta_2, 1)] \\
\downarrow a & & \downarrow a \\
[(h_0, h_1, h_2, \bar{\eta}_3, 1, \bar{\eta}_1)] & \xrightarrow{a_G?} & [(h_0, h_1, h_2, 1, \bar{\eta}_0, \bar{\eta}_2)]
\end{array}$$

where $\bar{\eta}_i = f_2(\eta_i)$. Since $a_G([\bar{\eta}_3, 1, \bar{\eta}_1]) = [(1, m_0(\bar{\eta}_1, \bar{\eta}_2, \bar{\eta}_3), \bar{\eta}_2)]$, f_2 preserves the 3-multiplications if and only if $m_0(\bar{\eta}_1, \bar{\eta}_2, \bar{\eta}_3) = \bar{\eta}_0$, i.e. if and only if the above diagram commutes. So 2b is also proven.

Translating the diagrams in item (2c) in groupoid language, we have

$$\begin{array}{ccc}
(J_l^{-1}(M \times_M H_0) \times_{G_0} G_1)/H_1 & \xrightarrow{\text{restriction of } a} & M \times_M H_0 \times_{M \times_M G_0} J_l^{-1}(M \times_M G_0) \\
\downarrow b_l^H \downarrow b_r^H & & \downarrow b_l^G \downarrow b_r^G \\
(H_1 \times_{G_0} G_1)/H_1 & \xrightarrow{f_2} & M \times_M H_0 \times_{M \times_M G_0} G_1
\end{array} \quad (20)$$

where J_l denotes the left moment map of X_2 or Y_2 to $G_0 \times_M G_0$ or $H_0 \times_M H_0$. Then the left diagram of (2c) is trivial since $b_l^{H,G} = id$ by the construction in Section 5. For the right diagram, we need to show that these two maps are the same: $[(\eta, 1)] \xrightarrow{a} (h_2, s_0(x), \bar{\eta}) \xrightarrow{b_r^G} (h_2, s_0(x), b_r^G(\bar{\eta}))$ and $[(\eta, 1)] \xrightarrow{b_r^H} [(b_r^H(\eta), 1)] \xrightarrow{f_2} (h_2, s_0(x), f_2(b_r^H(\eta)))$, where $x = d_1(h_2)$. That is, we need to show $b_r^G(f_2(\eta)) = f_2(b_r^H(\eta))$. Since $b_r = \varphi^{-1}$ is constructed by m 's as in Section 5, f_2 commutes with b_r 's. So we proved item (2c). $\square$

To establish the inverse argument, we fix again the notation: $\mathcal{G} \Rightarrow M$ is an SLie groupoid; $G_1 \Rightarrow G_0$ and $H_1 \Rightarrow H_0$ are two Lie groupoids presenting $\mathcal{G}$; X and Y are the 2-groupoids corresponding to G and H as constructed in Section 4.

Lemma 7.3. *If there is a Lie groupoid equivalence $H \rightarrow G$, then there is a Lie 2-groupoid equivalence $Y \rightarrow X$.*

Proof. Since both H and G present $\mathcal{G}$ which is an SLie groupoid over M , we are in the situation described in item 2 in Lemma 6.13, that is we have a 2-morphism a satisfying various commutative diagram as in item (2a), (2b), (2c). We take f_0 to be the isomorphism $M \cong M$, f_1 the map $H_0 \rightarrow G_0$, $f_2 : Y_2 \rightarrow X_2$ the map $\eta \mapsto [(\eta, 1)] \xrightarrow{a} (h_2, h_0, \bar{\eta}) \mapsto \bar{\eta}$. Since f_2 is made up by composition of smooth maps, it is a smooth map. Since $d_2 \times d_0$ is the moment map and a preserves the moment map, we have $h_i = d_i(\eta)$ for $i = 0, 2$. It is clear that f_0 and f_1 preserve the structure maps since they preserve $\bar{e}$, $\bar{s}$, $\bar{t}$ of $\mathcal{G} \Rightarrow M$. It is also clear that $d_i f_2(\eta) = f_1(h_i)$ for $i = 2, 0$ since $(h_2, h_0, \bar{\eta} = f_2(\eta)) \in H_0^2 \times_{G_0^2} X_2$. Since a preserves moment maps, $f_1(d_1(\eta)) = J_r([\eta, 1]) = J_r(h_2, h_0, \bar{\eta}) = d_1(\bar{\eta})$, where J_r is

the moment map to G_0 of the corresponding bibundles. Hence f_2 preserves the degeneracy maps.

For the face maps $s_0, s_1 : \square_1 \rightarrow \square_2$, we again look at the commutative diagram (20). Recalling $s_0(h) = b_l^{-1} \bar{e}_H(h)$, we have

$$f_2(s_0(h)) = pr_X a([(b_l^{-1} \bar{e}_H(h), 1)]) = b_l^{-1} \bar{e}_G f_1(h) = s_0 f_1(h),$$

where pr_X is the projection $H_0^2 \times_{G_0^2} X_2 \rightarrow X_2$ and similarly for s_1 using the diagram for b_r . Hence f_2 preserves the face maps.

The fact that f_2 preserves the 3-multiplications follows from the proof of item (2b) of Lemma 7.2.

Then the induced map $\phi : Y_2 \rightarrow \text{hom}(\partial\Delta^2, Y) \times_{\text{hom}(\partial\Delta^2, X)} X_2$ is $\eta \mapsto [(\eta, 1)] \xrightarrow{a} (h_2, h_0, \bar{\eta}) \mapsto (h_0, h_1, h_2, \bar{\eta})$, where $\bar{\eta} = f_2(\eta)$ and $h_i = d_i(\eta)$. As a composition of smooth maps ϕ is smooth. Moreover ϕ is injective since a is injective. For any $(h_0, h_1, h_2, \bar{\eta}) \in \text{hom}(\partial\Delta^2, Y) \times_{\text{hom}(\partial\Delta^2, X)} Y_2$, we have $(h_0, h_2, \bar{\eta}) \in H_0 \times_M H_0 \times_{G_0 \times_M G_0} X_2$. Then there is an η such that $[(\eta, 1)] = a^{-1}(h_0, h_2, \bar{\eta})$. Thus $\phi(\eta) = (h_0, d_1(\eta), h_2, \bar{\eta})$, which implies that $f_1(d_1(\eta)) = d_1(\bar{\eta}) = f_1(h_1)$. Therefore $(1, d_1(\eta), h_1) \in H_1$ and $d_i(\eta \cdot (1, d_1(\eta), h_1)) = h_i$, $i = 0, 1, 2$ since d_1 is the moment map to H_0 of the bibundle Y_2 . So $\phi(\eta \cdot (1, d_1(\eta), h_1)) = (\bar{\eta}, h_0, h_1, h_2)$ which shows the surjectivity. Therefore ϕ is an isomorphism. $\square$

The theorem is now proven, since we only have to consider the case that when (1-) Morita equivalence is given by Lie (2-) groupoid morphisms.

8 Comments on the inverse map

In this section, we prove that the axioms involving the inverse map in the definition of SLie groupoid are guaranteed by the axioms of multiplication and the identity. There is similar treatment of the antipode in hopfish algebras [TWZ]. In fact SLie groups are a geometric version of hopfish algebras.

Let $\mathcal{G} \Rightarrow M$ be an SLie groupoid, and $G := G_1 \xrightarrow[\mathbf{t}_G]{s_G} G_0$ a good groupoid presentation of $\mathcal{G}$ as we described in Section 3.1. So there is a map $e : M \rightarrow G_0$ presenting $\bar{e}$. We look at the following diagram,

$$\mathcal{G} \times_M \mathcal{G} \xrightarrow{m} \mathcal{G} \xleftarrow{\bar{e}} M,$$

and its corresponding groupoid picture,

$$\begin{array}{ccccc} G_1 \times_M G_1 & & G_1 & & M \\ \Downarrow & & \Downarrow & & \Downarrow \\ & \swarrow J_l & E_m & \searrow J_r & \\ G_0 \times_M G_0 & & G_0 & & M \\ & \swarrow J_l & E_{\bar{e}} & \searrow J_r & \\ & & & & \end{array} \quad (21)$$

where E_m and $E_{\bar{e}} = G_1 \times_{\mathbf{t}_G, G_0, e} M$ are bibundles presenting the multiplication m and identity $\bar{e}$ of $\mathcal{G}$ respectively. We can form a $G \times_M G$ left module $E_m \times_{G_0} E_{\bar{e}}/G$. Examining

the G action on $E_{\bar{e}}$, we see that the geometric quotient (corresponding to $\text{hom}_{\mathcal{A}}(\epsilon, \Delta)$ in the case of hopfish algebra),

$$(E_m \times_{G_0} E_{\bar{e}})/G = E_m \times_{J_r, G_0, e} M,$$

is still a manifold with a left $G_1 \times_M G_1$ action (which might not be free and proper). Therefore, we can view it as a left G module with the left action of the first copy of G and a right G^{op} module with the left action of the second copy of G . Here G^{op} is G with the opposite groupoid structure.

Lemma 8.1. *The bibundle above is a Morita equivalence from G to G^{op} with moment maps $pr_1 \circ J_l$ and $pr_2 \circ J_r$.*

Proof. Let $f_1 : \mathcal{G} \times_M \mathcal{G} \rightarrow \mathcal{G} \times_M \mathcal{G}$ be given by $(g_1, g_2) \mapsto (g_1 \cdot g_2, g_2)$, i.e. $f_1 = m \times pr_2$; let $f_2 : \mathcal{G} \times_M \mathcal{G} \rightarrow \mathcal{G} \times_M \mathcal{G}$ be given by $(g_1, g_2) \mapsto (g_1 \cdot g_2^{-1}, g_2)$. Since we have

$$(g_1, g_2) \xrightarrow{f_1} (g_1 g_2, g_2) \xrightarrow{f_2} ((g_1 g_2) g_2^{-1}, g_2) \sim (g_1, g_2),$$

and

$$(g_1, g_2) \xrightarrow{f_2} (g_1 g_2^{-1}, g_2) \xrightarrow{f_1} ((g_1 g_2^{-1}) g_2, g_2) \sim (g_1, g_2),$$

we have that $f_1 \circ f_2$ and $f_2 \circ f_1$ are isomorphic to id via 2-morphisms. Therefore f_1 is an isomorphism of stacks. Therefore $E_m \times_{pr_2 \circ J_l, G_0, \mathbf{t}_G} G_1$ presenting f_1 is a Morita bibundle from the Lie groupoid $G_1 \times_M G_1 \rightrightarrows G_0 \times_M G_0$ to $G_1 \times_M G_1 \rightrightarrows G_0 \times_M G_0$. Here the two moment maps are J_l (of E_m) and $J_r \times \mathbf{s}_G$. Therefore the left groupoid action of $G_1 \times_M G_1$ is principal on the bibundle $E_m \times_{pr_2 \circ J_l, G_0, \mathbf{t}_G} G_1$. The left action of the first copy of G_1 on E_m is the left action we need. Therefore, it is naturally free. To see that the desired action is also transitive, take (η_1, x) and (η_2, x) in $E_m \times_{J_r, G_0, e} M$ which project to the same point g_0 in G_0 by $pr_2 \circ J_l$. Then $(\eta_1, 1_{g_0})$ and $(\eta_2, 1_{g_0})$ are both in $E_m \times_{pr_2 \circ J_l, G_0, \mathbf{t}_G} G_1$ and both project to $(e(x), g_0) \in G_0 \times_M G_0$ via $J_r \times \mathbf{s}_G$. Therefore, there is an element $(\gamma_1, \gamma_2) \in G_1 \times_M G_1$ such that $(\gamma_1, \gamma_2) \cdot (\eta_1, 1_{g_0}) = (\eta_2, 1_{g_0})$. Therefore $\gamma_1 \cdot (\eta_1, x) = (\eta_2, x)$, i.e. the action is transitive fibre-wise.


The proof of the principality of the G^{op} action is similar (one considers $G_1 \times_{G_0} E_m$). $\square$

Remark 8.2. Another fibre product $E_m \times_{pr_2 \circ J_l, G_0, e} M$ is isomorphic to G_1 trivially via b_r . But the morphisms we use to construct the fibre product are different.

Notice that using the inverse operation, a G^{op} module is also a G module. In other words, the above lemma says that $E_m \times_{J_r, G_0, e} M$ is a Morita bibundle between G and G where the right G action is via the left action of the second of copy of $G \times_M G$ composed with the inverse. With this viewpoint, we have a stronger statement:

Lemma 8.3. *As Morita bibundles from G to G , $E_m \times_{J_r, G_0, e} M$ and E_i are isomorphic.*

This lemma is not at all surprising, if we use the view point of Lie 2-groupoids: $E_m \times_{J_r, G_0, e} M$ should be imagined as degenerate triangles (on the left)



Proof. We know from the property of E_i that $g \cdot g^{-1} \sim 1$, i.e. $((G_1 \times_{\mathbf{t}_G, G_0, J_l} E_i) \times_{\mathbf{s}_G \times J_r, G_0 \times_M G_0} E_m) / G_1 \times_M G_1$ is 2-isomorphic to the module $G_0 \times_{e \circ \mathbf{t}, G_0, \mathbf{t}_G} G_1$ presenting the map $e \circ \mathbf{t} : \mathcal{G} \rightarrow M \rightarrow \mathcal{G}$. We will first show that $E_m \times_{G_0} M$ also has this property.

Let $[(1, \eta_1, \eta_2)] \in (((G_1 \times_{G_0} (E_m \times_{J_r, G_0, e} M)) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1)$ (See (22)). We can suppose the first element to be 1 because the $G_1 \times_M G_1$ action on G_1 is simply right multiplication by the first factor. Moreover, examining the morphisms in the fibre product, we have $J_l(\eta_1) = J_l(\eta_2)$, and the right $G_1 \times_M G_1$ action is

$$(\gamma_0, \eta_1, \eta_2) \cdot (\gamma_1, \gamma_2) = (\gamma_0 \cdot \gamma_1, (1, \gamma_2^{-1}) \cdot \eta_1, (\gamma_1, \gamma_2)^{-1} \cdot \eta_2). \quad (23)$$

Since the right action of G_1 on E_m is principal (now viewing E_m as a bibundle from $G \times_M G$ to G), and $J_l(\eta_1) = J_l(\eta_2)$, there exists a unique element $\gamma \in G_1$ such that $\eta_1 \cdot \gamma = \eta_2$. Suppose $[(1, \eta_1, \eta_2)] = [(1, \tilde{\eta}_1, \tilde{\eta}_2)]$, then by (23), there exists $\gamma_2 \in G_1$ such that $\eta_1 = (1, \gamma_2^{-1}) \cdot \tilde{\eta}_1$ and $\eta_2 = (1, \gamma_2^{-1}) \cdot \tilde{\eta}_2$. Since the right action and left action on a bibundle commute, γ is independent of the choice of representative of $[(1, \eta_1, \eta_2)]$. Therefore, we have a well-defined map

$$\phi : ((G_1 \times_{G_0} (E_m \times_{J_r, G_0, e} M)) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1 \rightarrow G_0 \times_{e \circ \mathbf{t}, G_0, \mathbf{t}_G} G_1,$$

by

$$[(1_g, \eta_1, \eta_2)] \mapsto (g, \gamma).$$

Here (g, γ) is an element of $G_0 \times_{e \circ \mathbf{t}, G_0, \mathbf{t}_G} G_1$ since $\mathbf{t}_G(\gamma) = J_r(\eta_1)$. Moreover, since smoothness is a local property and all the moment maps are smooth and the groupoid action is principal, ϕ is naturally smooth. ϕ is an isomorphism following from principality of the right G_1 action on E_m . Moreover, it is not hard to check that ϕ is equivariant and commutes with the moment maps of the bibundles. Therefore,

$$((G_1 \times_{G_0} (E_m \times_{J_r, G_0, e} M)) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1 \cong G_0 \times_{e \circ \mathbf{t}, G_0, \mathbf{t}_G} G_1$$

as Morita bibundles. One proceeds similarly to prove the other symmetric isomorphism corresponding to $g^{-1} \cdot g \sim 1$.

Let φ be the composed isomorphism

$$((G_1 \times_{G_0} (E_m \times_{J_r, G_0, e} M)) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1 \rightarrow ((G_1 \times_{G_0} E_i) \times_{G_0 \times_M G_0} E_m) / G_1 \times_M G_1. \quad (24)$$

Suppose $\varphi([(1_g, \eta_1, \eta_2)]) =([(1_g, \tilde{\eta}_1, \tilde{\eta}_2)])$ (we can still assume that the first component is 1 because the $G_1 \times_M G_1$ action on both side is the right multiplication by the first copy; we can assume that they are 1 at the same point because φ commutes with the moment maps on the left leg). Examining the morphisms inside the fibre products, we have

$$pr_1 \circ J_l(\eta_2) = \mathbf{t}_G(1_g) = pr_1 \circ J_l(\tilde{\eta}_2) = g.$$

Since φ commutes with the moment maps on the right leg, we have

$$J_r(\eta_2) = J_r(\tilde{\eta}_2).$$

Similarly to the proof of Lemma 8.1, we can show that $G_1 \times_{\mathbf{s}_G, G_0, pr_1 \circ J_l} E_m$ is a Morita bibundle from $G \times_M G$ to $G \times_M G$. Then $(1_g, \eta_2)$ and $(1_g, \tilde{\eta}_2)$ are both in $G_1 \times_{\mathbf{s}_G, G_0, pr_1 \circ J_l} E_m$

and their images under the right moment map $\mathbf{s}_G \times J_r$ are both $(g, J_r(\eta_2))$. By principality of this left $G_1 \times_M G_1$ action, there is a unique $(\gamma_1, \gamma_2) \in G_1 \times_M G_1$ such that

$$(\gamma_1, \gamma_2) \cdot (1, \eta_2) = (1, \tilde{\eta}_2).$$

Therefore $\gamma_1 = 1$ and $(1, \gamma_2) \cdot \eta_2 = \tilde{\eta}_2$. This left $G_1 \times_M G_1$ action on E_m is exactly the left $G_1 \times_M G_1$ action on the second copy of E_m in (24). Using this γ_2 , we have

$$(1, \tilde{\eta}_1, \tilde{\eta}_2) \cdot (1, \gamma_2) = (1, \gamma_2^{-1}) \cdot (1, \tilde{\eta}_1, \tilde{\eta}_2) = (1, \eta'_1, \eta_2).$$

Therefore the isomorphism

$$\varphi : [(1_g, \eta_1, \eta_2)] \mapsto [(1_g, \eta'_1, \eta_2)]$$

induces a map $\psi : E_m \times_{G_0} M \rightarrow E_i$ by $\eta_1 \mapsto \eta'_1$. It's routine to check ψ is an isomorphism of Morita bibundles. $\square$

As in this Lemma, we have seen that the 2-identities satisfied by E_i are actually naturally satisfied by $E_m \times_{J_r, G_0, e} M$. Notice that for the first part of the proof, we didn't use any information involving the inverse map. Our conclusion is that the inverse map represented by E_i can be replaced by $E_m \times_{J_r, G_0, e} M$ without any further conditions (not even on the 2-morphisms). In fact, the natural 2-morphisms coming along with bibundle $E_m \times_{J_r, G_0, e} M$ naturally go well with other 2-morphisms a 's and b 's.

Proposition 8.4. *An SLie groupoid $\mathcal{G}$ can be equally defined by all the axioms except the axioms involving inverses and the additional condition that $E_m \times_{J_r, G_0, e} M$ is a Morita bibundle from G to G^{op} for some good presentation G of $\mathcal{G}$.*

Proof. It is clear from the Lemma 8.3 that the existence of the inverse map guarantees that the bibundle $E_m \times_{J_r, G_0, e} M$ is a Morita bibundle from G to G^{op} for a good presentation G of $\mathcal{G}$.

On the other hand, if $E_m \times_{J_r, G_0, e} M$ is a Morita bibundle from G to G^{op} for some presentation G of $\mathcal{G}$, then we construct the inverse map $i : \mathcal{G} \rightarrow \mathcal{G}$ by this bibundle. Because of the nice properties of $E_m \times_{J_r, G_0, e} M$ that we have proven in the first half of Lemma 8.3, this newly defined inverse map satisfies all the axioms that the inverse map satisfies. $\square$

Remark 8.5. This theorem holds also for W-groupoids and the proof is similar. Then one can see that the new definition of SLie group modulo 2-morphisms is analogous to the definition of hopfish algebra.

Sometimes the inverse of an SLie groupoid is given by a specific groupoid isomorphism $i : G \rightarrow G$ on some presentation (for example $\mathcal{G}(A)$ and $\mathcal{H}(A)$ in [TZ06a] and (quasi-)Hopf algebras as the algebra counter-part).

Lemma 8.6. *The inverse map of an SLie groupoid $\mathcal{G}$ is given by a groupoid isomorphism $i : G \rightarrow G$ for some presentation G if and only if on this presentation $E_m \times_{J_r, G_0, e} M$ is a trivial right G principal bundle over G_0 .*

Proof. It follows from Lemma 8.3 and the fact that the inverse is given by a morphism $i : G \rightarrow G$ if only if the bibundle E_i is trivial. $\square$

References

- [AGV72] M. Artin, A. Grothendieck, and J. L. Verdier. *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*. Springer-Verlag, Berlin, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 269.
- [AM84] Rui Almeida and Pierre Molino. Suites d’Atiyah, feuilletages et quantification géométrique. *Université des Sciences et Techniques de Languedoc, Montpellier, Séminaire de géométrie différentielle*, pages 39–59, 1984.
- [AM85] Rui Almeida and Pierre Molino. Suites d’Atiyah et feuilletages transversalement complets. *C. R. Acad. Sci. Paris Sér. I Math.*, 300(1):13–15, 1985.
- [BCSS] John C. Baez, Alissa S. Crans, Danny Stevenson, and Urs Schreiber. From Loop Groups to 2-Groups, math.QA/0504123.
- [BL04] John C. Baez and Aaron D. Lauda. Higher-dimensional algebra. V. 2-groups. *Theory Appl. Categ.*, 12:423–491 (electronic), 2004.
- [BM94] J.-L. Brylinski and D. A. McLaughlin. The geometry of degree-four characteristic classes and of line bundles on loop spaces. I. *Duke Math. J.*, 75(3):603–638, 1994.
- [BS76] Ronald Brown and Christopher B. Spencer. Double groupoids and crossed modules. *Cahiers Topologie Géom. Différentielle*, 17(4):343–362, 1976.
- [BX] Kai Behrend and Ping Xu. Differentiable Stacks and Gerbes, math.DG/0605694.
- [BZ] Henrique Bursztyn and Chenchang Zhu. Morita equivalence of weinstein groupoids. in preparation.
- [CdSW99] Ana Cannas da Silva and Alan Weinstein. *Geometric models for noncommutative algebras*, volume 10 of *Berkeley Mathematics Lecture Notes*. American Mathematical Society, Providence, RI, 1999.
- [Dus79] J. Duskin. Higher-dimensional torsors and the cohomology of topoi: the abelian theory. In *Applications of sheaves (Proc. Res. Sympos. Appl. Sheaf Theory to Logic, Algebra and Anal., Univ. Durham, Durham, 1977)*, volume 753 of *Lecture Notes in Math.*, pages 255–279. Springer, Berlin, 1979.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [Gle82] Paul G. Glenn. Realization of cohomology classes in arbitrary exact categories. *J. Pure Appl. Algebra*, 25(1):33–105, 1982.
- [Hen] Andre Henriques. Integrating L-infinity algebras, math.AT/0603563.

- [HZ] André Henriques and Chenchang Zhu. Stacky lie groupoids and lie 2-groupoids. in preparation.
- [May92] J. Peter May. *Simplicial objects in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992. Reprint of the 1967 original.
- [Met] David Metzler. Topological and smooth stacks, math.DG/0306176.
- [Moe02] Ieke Moerdijk. Orbifolds as groupoids: an introduction. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 205–222. Amer. Math. Soc., Providence, RI, 2002.
- [Mrč] J Mrčun. *Stablility and invariants of Hilsum-Skandalis maps*. Dissertation, Utrecht University, Utrecht, 1996.
- [Noo] Behrang Noohi. Notes on 2-groupoids, 2-groups and crossed-modules, math.CT/0512106.
- [Pro96] Dorette A. Pronk. Etendues and stacks as bicategories of fractions. *Compositio Math.*, 102(3):243–303, 1996.
- [ST04] Stephan Stolz and Peter Teichner. What is an elliptic object? In *Topology, geometry and quantum field theory*, volume 308 of *London Math. Soc. Lecture Note Ser.*, pages 247–343. Cambridge Univ. Press, Cambridge, 2004.
- [Sto96] Stephan Stolz. A conjecture concerning positive Ricci curvature and the Witten genus. *Math. Ann.*, 304(4):785–800, 1996.
- [TWZ] Xiang Tang, Alan Weinstein, and Chenchang Zhu. Hopfish algebras. math.QA/0510421, to appear in Pacific Journal of Mathematics.
- [TZ06a] Hsian-Hua Tseng and Chenchang Zhu. Integrating Lie algebroids via stacks. *Compos. Math.*, 142(1):251–270, 2006.
- [TZ06b] Hsian-Hua Tseng and Chenchang Zhu. Integrating Poisson manifolds via stacks. *Travaux mathématique*, 15:285–297, 2006.
- [Vis02] Angelo Vistoli. Topological stacks. *a talk given at MSRI*, (available at <http://www.msri.org/publications/ln/msri/2002/introstacks/vistoli/1/index.html>), 2002.
- [Š] Pavol Ševera. Some title containing the words “homotopy” and “symplectic”, e.g. this one, math.sg/0105080.
- [Wei83] Alan Weinstein. The local structure of Poisson manifolds. *J. Differential Geom.*, 18(3):523–557, 1983.
- [WX91] Alan Weinstein and Ping Xu. Extensions of symplectic groupoids and quantization. *J. Reine Angew. Math.*, 417:159–189, 1991.
- [Zhu] Chenchang Zhu. Lie’s II theorem of Lie algebroids via stacky Lie groupoids. in preparation.